SCIENCE.

EDITORIAL COMMITTEE: S. NEWCOMB, Mathematics; R. S. WOODWARD, Mechanics; E. C. PICKERING, Astronomy; T. C. MENDENHALL, Physics; R. H. THURSTON, Engineering; IRA REMSEN, Chemistry; JOSEPH LE CONTE, Geology; W. M. DAVIS, Physiography; O. C. MARSH, Paleontology; W. K. BROOKS, Invertebrate Zoölogy; C. HART MERRIAM, Vertebrate Zoölogy; N. L. BRITTON, Botany; HENRY F. OSBORN, General Biology; H. P. BOWDITCH, Physiology; J. S. BILLINGS, Hygiene; J. MCKEEN CATTELL, Psychology; DANIEL G. BRINTON, J. W. POWELL, Anthropology.

FRIDAY, MARCH 22, 1895.

CONTENTS:

The Fundamental Difference between Animals and The Best Order of Topics in a Two-years' Course of Anatomy in a Medical School: Frederic Henry GERRISH312 Current Notes on Physiography (IV.): W. M. Annual Reception of the New York Academy: HENRY F. OSBORN......321 An International Scientific Catalogue and Congress: HORATIO HALE. Brinton's Primer of Mayan Hieroglyphics: FRED-ERICK STARR. Yeo's Steam Engine: R. H. THURSTON. Life of Dean Buckland: A. S.

PACKARD. Geology.

Notes and News :-

 Scientific Journals
 335

 New Books
 336

MSS. intended for publication and books, etc., intended for review should be sent to the responsible editor, Prof. J. McKeen Cattell, Garrison on Hudson, N. Y. Subscriptions and advertisements should be sent to Science, 41 N. Queen St., Lancaster, Pa., or 41 East 49th St., New York.

ARGON.

THE plain facts concerning argon are these: For some time past Lord Rayleigh has been engaged on refined work involving the weighing of various gases. Last year he found that the nitrogen obtained from the air is a little heavier than that made from definite chemical compounds. This led him to further experiments and, at the same

time, Professor W. Ramsay, of University College, London, also undertook experiments with the object of explaining, if possible, the discrepancy. The general method of work consisted in passing air, first through substances that have the power to remove those constituents that are present in small quantities, such as water vapor, carbonicacid gas, etc., then through a heated tube containing copper. The oxygen of the air unites with the heated copper, and what has hitherto been regarded as nitrogen remains uncombined. This 'atmospheric nitrogen' was subsequently treated in three different ways for the purpose of removing the nitrogen from it.

- (1) It was drawn through clay pipes in the hope that, if the gas is a mixture, one of the constituents would pass through the porous material more easily than the other, and at least a partial separation be thus effected. While something was accomplished in this way, the experiment was on the whole unsatisfactory.
- (2) The 'atmospheric nitrogen' was mixed with oxygen in a vessel containing caustic alkali, and electric sparks were passed through the mixture. Under these circumstances the oxygen united with nitrogen and formed a compound which is soluble in alkali. After no further absorption of nitrogen could be effected by sparking, any unchanged oxygen present was removed, and there was then found a residue



1	(1	2

Science 1 (12), 309-336.

ARTICLE TOOLS http://science.sciencemag.org/content/1/12.citation

PERMISSIONS http://www.sciencemag.org/help/reprints-and-permissions

Use of this article is subject to the Terms of Service