

genesis in *Drosophila* (Sobels). Harm discussed repair of lesions by enzymes and the idea that repair, to be effective, must occur before DNA synthesis; this is becoming an increasingly attractive explanation for the various findings with *Escherichia*, *Paramecium*, and *Drosophila*.

The influence of dose rate and dose fractionation on mutation induction by ionizing radiation was discussed for the mouse (W. L. Russell), silkworm (Y. Tazima and S. Kondo), and *Drosophila* (H. J. Muller, I. I. Oster and S. Zimmering, and C. E. Purdom). The existence of dose rate and fractionation effects is now well established for the mouse and the silkworm but the data for *Drosophila* are still equivocal. Some of the effects in the mouse and silkworm can be explained best by hypotheses involving repair of pre-mutational damage. Other effects require hypotheses postulating some form of selection, such as differential cell killing.

Concerning the changes in sensitivity during gametogenesis and early cleavage stages, a number of striking parallels are found between diverse organisms. A rather full description was given by R. C. von Borstel and W. St. Amand of the changes in sensitivity from early oögenesis to early cleavage in *Habrobracon*. F. E. Würigler studied in detail the changes in sensitivity of *Drosophila* eggs during the first few cleavages. The *Habrobracon* and *Drosophila* studies agree in showing a peak of sensitivity just after meiosis in fertilized eggs and a periodic rise and fall in sensitivity corresponding to the first few cleavage mitoses.

The sensitivity of sperm and of various stages of spermatogenesis in *Drosophila* were discussed. R. Sävhaugen summarized her work on the variation of sensitivity to induction of various types of genetic effects during spermatogenesis. J. Mossige considered in some detail the factors responsible for the difference in sensitivity between various batches of mature sperm. H. Traut analyzed the basis for variation in the form of the dose curves for recessive lethals in sperm of different origins. D. L. Lindsley pointed out the relations between the chromosome constitution of the sperm and its sensitivity to inactivation by radiation. B. P. Kaufman and H. Gay discussed electron microscopical and cytochemical studies of spermatogenesis.

Work by L. B. Russell on the sensitivity of various stages in gametogenesis and in the early development of the

mouse zygote is now extensive enough to allow a number of comparisons with the studies on *Drosophila* and *Habrobracon*. These comparisons result in several striking parallels. G. E. Magni discussed the major changes in sensitivity that occur during meiosis in yeast and compared the sensitivity of mitosis and meiosis in this organism, both in regard to spontaneous and radiation-induced mutation.

The symposium was organized by F. H. Sobels who assembled a group of approximately 100 major investigators in the field. There were six half-day sessions and an evening session with a total of 24 invited papers. There was adequate time for discussion, and the discussion was lively. The conference was well organized and succeeded admirably in its purpose of defining the present status of research in this general area in the presence of most of the active workers. The invited papers and recorded discussion will be published by Pergamon Press.

Important advances have been made in understanding the mechanisms by which the initial radiation damage to the chromosomes is converted to chromosome aberrations or mutations, and it is becoming possible to apply this knowledge to interpreting variations in radiation sensitivity. The existence of dose rate and dose fractionation effects in the induction of specific locus mutations can no longer be doubted for either the mouse or the silkworm although the situation remains equivocal for *Drosophila*. Work in this area is turning now to analyses of the reasons for these effects. Many data have been accumulated on variations in radiation sensitivity during gametogenesis in various organisms, and some striking similarities have been found between quite diverse organisms. Though a number of more or less plausible explanations have been offered for the variations in sensitivity, general agreement has not yet been reached; the major advances in this area have been more descriptive than analytical.

R. F. KIMBALL


Biology Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee

Forthcoming Events

January

23-26. American Assoc. of Physics Teachers, New York, N.Y. (R. P. Winch, Williams College, Williamstown, Mass.)
23-26. American Group Psychotherapy

ACCESSORIES INCREASE CAPABILITIES OF THE MODEL 350 SPECTROPHOTOMETER



Highly versatile as a basic analytical instrument, the Perkin-Elmer Model 350 UV-VIS-NIR Spectrophotometer has even more impressive capabilities with accessories to perform extra functions. Here are some of the auxiliaries that are available: **Controlled-Temperature Cell Mount** (illustrated), for Perkin-Elmer Cylindrical Sample Cells, maintains any specified temperature from 0°C to 100°C within 0.5°C. Helps determine the kinetics of reactions at various temperatures.

Standard Time Drive Accessory—records transmittance or absorbance against time at any of 11 different speeds, for precise one-point kinetic studies.

Repetitive Scan Accessory enables the operator to re-scan any selected segment of an absorption spectrum automatically at regular time intervals, to analyze general chemical reaction changes.

Linear Wavelength Accessories provide UV, VIS or NIR linear wavelength readout on the standard preprinted Model 350 chart, where this mode of recording is desired.

Auxiliary Recorder Readout produces additional spectra for any desired purpose: enables operator to stretch or compress spectra, by chart speed variations. Useful with Linear Wavelength Accessories, above.

Other accessories include Potassium Bromide Disc Mount, Short Fixed-Path Liquid Cell, and Variable-Path Liquid Cell. For complete details on Model 350 accessories, write to Instrument Division, Perkin-Elmer Corporation, 910 Main Avenue, Norwalk, Connecticut.

PERKIN-ELMER

SCIENCE, VOL. 139