

Comment. It is fully realized that hardly any of the *Mycoplasma* species hitherto described do in fact strictly meet the requirements that are formulated in paragraph 6, a. It is appreciated, moreover, that self-evident though these requirements are in principle, it will prove difficult or even impossible for any single worker or group of workers to satisfy the demands. However, it is hoped that the gradual creation of a network of reference laboratories may help to ameliorate the situation in this respect and steps are being taken toward this end.

At any rate, rather than compromising too much with the above requirements, it would be wise policy to restrain one's taxonomic efforts and to publish any new isolates merely under their catalog designations, thus providing a useful and necessary means of reference until it is possible to provide a reasonably adequate description.

Subcommittee on the Taxonomy of Mycoplasmatales:

D. G. FF. EDWARD, *Chairman*
Wellcome Research Laboratories,
Beckenham, Kent, England

E. A. FREUNDT, *Secretary*
University of Aarhus,
Aarhus, Denmark

R. M. CHANOCK
National Institutes of Health,
Bethesda, Maryland

J. FABRICANT
New York State Veterinary College,
Cornell University, Ithaca

L. HAYFLICK
Wistar Institute,
Philadelphia, Pennsylvania

R. M. LEMCKE
Lister Institute of Preventive
Medicine, London, England

S. RAZIN
Hebrew University, Hadassah Medical
School, Jerusalem, Israel

N. L. SOMERSON
Children's Hospital, Columbus, Ohio

RUTH G. WITTLER
Walter Reed Army Institute of
Research, Washington, D.C.

References and Notes

1. Subcommittee on the Taxonomy of Mycoplasmatales, *Int. J. System. Bacteriol.*, in press.
2. D. G. ff. Edward and E. A. Freundt, *J. Gen. Microbiol.* 14, 197 (1956).
3. Editorial Board: International Code of Nomenclature of Bacteria, *Int. J. System. Bacteriol.* 16, 459 (1966).
4. A. B. Sabin, *Bacteriol. Rev.* 5, 1 (1941).
5. L. Hayflick and R. M. Chanock, *ibid.* 29, 185 (1965).
6. H. C. Neimark and J. J. Pene, *Proc. Soc. Exp. Biol. Med.* 118, 517 (1965).
7. D. G. ff. Edward and E. A. Freundt, *Int. J. System. Bacteriol.*, in press.
8. D. G. ff. Edward, *Int. Bull. Bacteriol. Nomencl.* 5, 85 (1955).

9. M. Rogul, Z. A. McGee, R. G. Wittler, S. Falkow, *J. Bacteriol.* 90, 1200 (1965).
10. P. R. Reich, N. L. Somerson, J. A. Rose, S. M. Weissman, *ibid.* 91, 153 (1966).
11. P. R. Reich, N. L. Somerson, C. J. Hybner, R. M. Chanock, S. M. Weissman, *J. Bacteriol.* 92, 302 (1966).
12. N. L. Somerson, P. R. Reich, B. E. Walls, R. M. Chanock, S. M. Weissman, *ibid.*, p. 311.
13. Cultures may be sent to the American Type Culture Collection (12301 Parklawn Drive, Rockville, Maryland 20852) or to the National Collection of Type Cultures (Central Public Health Laboratory, Colindale Avenue, London, N.W.9, England).
14. *International Code of Nomenclature of Bacteria and Viruses. Bacteriological Code*, Editorial Board of the International Committee on Bacteriological Nomenclature, Eds. (Iowa State College Press, Ames, 1958).

13 March 1967

Sexual Reproduction in *Histoplasma capsulatum*

Recent successes (1) in stimulating sexual reproduction among the dermatophytes and related fungi inspired a search for the perfect state of fungi that cause systemic disease in man. One of the organisms under study was *Histoplasma capsulatum*, the etiologic agent of histoplasmosis, a pulmonary disease of global importance. Twenty-nine isolates of this mold, recovered from soil, bats, or humans, were grown singly or in combination on small pieces of sterilized chicken feathers or horse hairs placed on plates of moistened, sterilized soil.

Cleistothecia filled with asci and ascospores were formed by two of the isolates. With the aid of a micro-manipulator, single ascospore cultures were obtained, and the organism was found to be homothallic.

The two cultures that developed the cleistothecia (H-2 and H-8) had been isolated from soil collected under a starling (*Sturnus vulgaris*) roost in Illinois (H-2) and from a case of histoplasmosis (H-8) in Puerto Rico.

The morphological characteristics of the cleistothecia and their asci were typical of the genus *Gymnoascus* of the family Gymnoascaceae. The perfect state of *Histoplasma capsulatum* is being studied.

LIBERO AJELLO
SHU-LAN CHENG

National Communicable Disease Center,
Atlanta, Georgia 30333

References

1. P. Stockdale, *Sabouraudia* 1, 41 (1961); C. O. Dawson and J. C. Gentles, *ibid.*, p. 49; L. Ajello, *ibid.*, p. 173; P. Stockdale, D. W. R. Mackenzie, P. K. C. Austwick, *ibid.* 4, 112 (1965).

6 February 1967

Medial Superior Olive and Sound Localization

Harrison and Irving (1) show that some animals which are capable of localizing a sound source do not possess a medial superior olive. On this basis, the authors rule out the idea that the medial superior olive is essential for sound localization. However, the electrophysiological data which they cite (2), together with some recent behavioral data not available to them (3), suggest an alternative interpretation of the presence or absence of the medial superior olive in specific mammals.

In order to understand the contribution of the medial superior olive to sound localization, it is necessary to distinguish between the two potential cues in the stimulation reaching the ears, both of which vary with the azimuth of the source of a brief sound (4). The first of these potential cues is the difference in the time of arrival of the wave front at the two ears. This time difference, Δt , depends directly on the distance between the ears and inversely with the speed of sound in the conducting medium. Animals with small heads and aquatic animals with even moderately large heads are virtually deprived of this potential cue since the interaural distance is small or the speed of sound is too great.

The second potential cue for the localization of the source of a brief sound is the differences in the frequency spectrum of the stimulation reaching the two ears (5). The spectrum difference, $\Delta(f)$, depends on the effectiveness of the sound shadow produced by the head and pinna. No animal is completely deprived of this potential cue, but since the effectiveness of the sound shadow decreases with a decrease in the proportion of high frequencies in the stimulation, animals (such as man) which are relatively insensitive to high frequencies are not exposed to a wide range in $\Delta(f)$. Thus Δt is the more dramatic cue for azimuth in animals with wide-set ears while $\Delta(f)$ is the more dramatic cue in animals with close-set ears, an aquatic habitat, or sensitivity to high frequencies.

The analysis of the Δt cue and the analysis of the $\Delta(f)$ cue are accomplished by different structures in the auditory system. We have isolated Δt and $\Delta(f)$ by delivering clicks through headphones worn by experimental ani-

Downloaded from http://science.sciencemag.org/ on January 20, 2019

mals (3). We observed that, first, surgical transection of the afferent tracts to the superior olive permanently abolishes Δt discrimination throughout the range which occurs in natural sound localization. Second, transection of lemniscal pathways from hindbrain to auditory cortex also abolishes Δt discrimination. Third, the same lesions which abolish Δt discrimination have little effect either on tasks requiring discriminations within the generic class $\Delta(f)$ or on sound localization itself (6).

The first result, together with electrophysiological data showing that Δt analysis takes place in the medial superior olive but not in the lateral superior olive, suggests that the role of the medial superior olive in sound localization is the analysis of Δt . The second result shows that the product of the analysis in the medial superior olive must be available to the forebrain in order for the animal to discriminate Δt . The third result shows that an animal deprived of Δt analysis is not necessarily incapable of sound localization because $\Delta(f)$ analysis is still available.

If we assume, as do Harrison and Irving, that the medial superior olive is homologous in all mammals and, further, that in the animals where it is present it is also similar in function, we conclude that animals without a medial superior olive may be insensitive to Δt and yet still capable of sound localization on the basis of $\Delta(f)$.

Finally, we are persuaded that the correlation between the size of the abducens nucleus and the size of the medial superior olive discovered by Harrison and Irving can hardly be a matter of chance (7) and, further, that the authors are probably correct in concluding that the visual system makes use of the Δt analysis provided by the medial superior olive. The size of the medial superior olive in auditory systems sensitive to Δt may indeed depend on its contribution to reflexive eye, ear, and head movements as Harrison and Irving argue, but this function is in addition to, not instead of, its contribution to the higher order instrumental activities generated by the forebrain.

BRUCE MASTERTON

Department of Psychology,
Vanderbilt University,
Nashville, Tennessee 37203

IRVING T. DIAMOND

Department of Psychology,
Duke University,
Durham, North Carolina 27706

References and Notes

1. J. M. Harrison and R. Irving, *Science* **154**, 738 (1966).
2. In addition to the references cited by Harrison and Irving, see M. R. Rosenzweig, *J. Comp. Physiol. Psychol.* **47**, 269 (1954); M. R. Rosenzweig and E. J. Wyers, *ibid.* **48**, 426 (1955); M. R. Rosenzweig and A. H. Amon, *Experientia* **11**, 498 (1955); M. R. Rosenzweig and D. Sutton, *J. Neurophysiol.* **21**, 17 (1958).
3. B. Masterton, J. A. Jane, I. T. Diamond, *J. Neurophysiol.*, in press. The technique employed is similar to that described in R. B. Masterton and I. T. Diamond, *ibid.* **27**, 15 (1964) except that the discrimination tests of Δt and $\Delta(f)$ were performed after lesions were placed in the trapezoid body or lateral lemniscus.
4. It is essential to restrict the discussion to brief sounds since any other kind can be easily localized by head movements and require no special analytic apparatus.
5. The second cue is usually called an intensity difference. This term was derived from research on localization of pure tones and, for brief sounds, is a bit misleading. Frequency spectrum difference is a more comprehensive term which includes intensity differences as special cases.
6. G. C. Nauman, thesis, University of Chicago (1958).
7. After reading Harrison and Irving's report, we counted the cells in the homologous nuclei of tree shrew (*Tupaia glis*), which has a retina containing only cones, and bushbaby (*Galago senegalensis*), which has a retina containing only rods. Although our cell counts place these animals intermediate to the two regression lines reported by Harrison and Irving, neither is so far off as to deny the general relation.
8. Supported by NIH grants NB-5251 and M-4849.

13 January 1967

Our work (1) on the nuclei of the superior olivary complex was essentially an anatomical study of the size of the medial superior olivary nucleus and of the relation between the size of the medial superior olivary nucleus and the size of the 6th nucleus. Since the medial superior olivary nucleus is absent in some animals, mammals can be divided into two groups—those with




Fig. 1. Schematic horizontal section through the acoustic system of the rat. The principal ascending nerve fibers which run in the ventral acoustic stria are shown. AN, Acoustic nerve; LF, large fibers of trapezoid body; LL, lateral lemniscus; LSO, lateral superior olivary nucleus; LTB, lateral nucleus of trapezoid body; MCP, middle cerebellar peduncle; MSO, medial superior olivary nucleus; MTB, medial nucleus of trapezoid body; P, pyramidal tract; PVC, posterior ventral cochlear nucleus; SF, small fibers of the trapezoid body; SPN, superior paraolivary nucleus; ST, spinal trigeminal tract; I, II, III, regions of the cochlear nucleus.

and those without the medial superior olive. That is, two types of auditory systems can be distinguished. These anatomical data tell nothing about the functional (behavioral and physiological) properties of these structures, except that in those animals in which the medial superior olive is missing (bats, for example) localization and other auditory behavior must depend upon other structures. Masterton and Diamond point out that auditory localization may be based on time, Δt , and intensity, $\Delta(f)$, differences of the stimulus at the two ears, and they suggest that one of these, the Δt discrimination, depends upon the medial superior olivary nucleus.

One way to investigate the role of the various nuclei of the superior olivary complex in behavior is to conduct the type of experiments reported by Masterton and Diamond. The problem of determining the role of these nuclei in behavior by the lesion method, however, is exceedingly complex because of the structural arrangements of the nerve fiber bundles and the nuclei of the ventral acoustic system of the medulla (2). Figure 1 shows some of the principal ascending bundles of axons which run in the ventral acoustic stria. Sectioning the afferent nerve fibers to the superior olivary complex will interrupt a number of ipsilateral and contralateral pathways (including that to the medial superior olive). Therefore, structural interpretations of the behavioral effects of such lesions are hazardous. Sectioning the afferent tracts to the superior olivary complex abolishes Δt discrimination, but because of the foregoing considerations many interpretations of such lesions are possible. The interpretation by Masterton and Diamond may be correct, but it could just as well be incorrect.

J. M. HARRISON
M. D. BEECHER

Psychological Laboratory,
Boston University,
Boston, Massachusetts 02215

References and Notes

1. J. M. Harrison and R. Irving, *Science* **154**, 738 (1966). Further information on the anatomical correlations will be found in R. Irving and J. M. Harrison, *J. Comp. Neurol.*, in press.
2. A number of papers on the ascending connections of the cochlear nucleus have recently appeared; the following are the most relevant to the present discussion: J. M. Harrison and R. Irving, *J. Comp. Neurol.* **126**, 51 (1966); J. M. Harrison, *Anat. Record* **154**, 463 (1966); R. Irving, *ibid.*, p. 466; T. P. S. Powell and S. D. Erulkar, *J. Anat.* **96**, 249 (1962); W. A. Stotler, *J. Comp. Neurol.* **98**, 401 (1953); B. Warr, *Exp. Neurol.* **14**, 453 (1966).

14 February 1967

Medial Superior Olive and Sound Localization

Bruce Masterton, Irving T. Diamond, J. M. Harrison and M. D. Beecher

Science **155** (3770), 1696-1697.
DOI: 10.1126/science.155.3770.1696-a

ARTICLE TOOLS <http://science.sciencemag.org/content/155/3770/1696.2.citation>

PERMISSIONS <http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)

Science (print ISSN 0036-8075; online ISSN 1095-9203) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. 2017 © The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. The title *Science* is a registered trademark of AAAS.