



The CHIME radio telescope consists of four fixed 20- by 100-m semicylinders. Along the axis of each cylinder, 256 dual-polarization feeds will be installed that will provide sensitivity in the range of 400 to 800 MHz. The full combination of all the feeds will be done by a correlator that will be far larger than any previously built and will provide over 200 square degrees of instantaneous sky coverage. This, together with a back-end burst searching instrument that will detect short radio signals in real time, will enable CHIME to detect many FRBs per day.

The CHIME (Canadian Hydrogen Intensity Mapping Experiment) radio telescope, currently being built in Penticton, British Columbia (see the photo), will be several orders more sensitive than the Caltech instrument and, depending on the true brightness distribution, could detect dozens of FRBs per day, even in the conventional, nonevolving Euclidean scenario. CHIME should thus blow the field open, enabling a wide variety of different FRB statistical analyses and a large number of localization opportunities. The CHIME FRB project is expected to come online in the latter half of 2017 and is eagerly awaited by the community.

Regardless of the brightness distribution, one thing is sure: FRBs are proving themselves to be truly useful probes on the extragalactic universe, hence providing a great boon—not just a boom—for all of astronomy. ■

REFERENCES

1. D. R. Lorimer *et al.*, *Science* **318**, 777 (2007).
2. L. Spitler *et al.*, *Nature* **531**, 202 (2016).
3. V. Ravi *et al.*, *Science* **354**, 1249 (2016).
4. H. K. Vedantham, V. Ravi, G. Hallinan, R. M. Shannon, <https://arxiv.org/abs/1606.06795> (2016).
5. N. Oppermann, L. Connor, U. Pen, *Mon. Not. R. Astron. Soc.* **461**, 984 (2016).

10.1126/science.aag1600

NEUROSCIENCE

Why does time seem to fly when we're having fun?

Increased dopamine may cause our internal clock to slow down, creating a sense of speeding time

By Patrick Simen¹ and Matthew Matell²

Animals use the neurotransmitter dopamine to encode the relationship between their responses and reward. Reinforcement learning theory (1) successfully explains the role of phasic bursts of dopamine in terms of future reward maximization. Yet, dopamine clearly plays other roles in shaping behavior that have no obvious relationship to reinforcement learning, including modulating the rate at which our subjective sense of time grows in real time. On page 1273 of this issue, Soares *et al.* (2) closely examine the role of dopamine in mice performing a task in which they keep track of the time between two events and make decisions about this temporal duration. The results suggest the need to reassess the leading theory of dopamine function in timing—the dopamine clock hypothesis (3). They may also help explain empirical phenomena that challenge the reinforcement learning account of dopamine function.

Reinforcement learning theory posits that reward prediction errors inform animals about which behaviors to engage in so as to maximize future reward. Time is the key factor in the experiments that have linked reinforcement learning to dopamine (4). These experiments demonstrate that behavior is shaped by bursts of activity (“phasic activity”) in midbrain dopamine-secreting neurons. When a reward occurs at an unexpected time, a positive reward prediction error is generated (increased dopamine release signals that “things are better than expected”). Consequently, communication from cortical neurons to neurons in the striatum is altered to change behavior in ways that maximize reward. Conversely, a negative reward prediction error signaled by a decrease in dopamine release occurs when a reward is not provided when expected. In addition, if delivery of a reward reliably occurs at the predicted time, it produces no burst of dopamine release at that moment,

but instead begins to produce a phasic dopamine signal in response to the earliest reward-predicting cue. Thus, future choices can be made so as to obtain the cue, and thereafter the reward. Eventually this process yields a temporally extended chain of cues and behaviors terminating in reward. However, time itself has typically been of less interest to reinforcement learning-dopamine researchers than the mechanisms that link these elements into chains of expected sensory inputs and planned motor outputs.

Yet, dopamine plays a key role in interval timing. The dopamine clock hypothesis holds that increased dopamine release speeds up an animal's subjective sense of time—its internal clock. For example, rats treated with amphetamine, which enhances dopamine release, respond earlier than when they are tested without the drug. Curiously, a simple prediction of the dopamine clock hypothesis would seem to be that time doesn't fly, but rather crawls, when you're having fun. Unexpectedly pleasurable events boost dopamine release, which should cause your internal clock to run faster. Your subjective sense of time in that case grows faster than time itself, so that short intervals seem longer than they are. The dopamine clock hypothesis accounts for this counterintuitive prediction by an additional assumption about attention: When things are good, attention to time is reduced, such that intervals seem shorter than they are (5).

To clarify the role of dopamine in interval timing, Soares *et al.* investigated midbrain dopamine neuron activity in the substantia nigra pars compacta (SNc) of mice performing a timing task. They presented mice with two brief tones, and trained them to classify the interval between the tones as shorter or longer than a standard criterion. They then observed calcium influx into dopaminergic SNc neurons, which signals activity. Consistent with standard reinforcement learning theory, the authors observed bursts of activity in dopamine-synthesizing neurons that were locked to the second tone, reflecting the probability of an upcoming reward. This probability was greatest when the in-

¹Department of Neuroscience, Oberlin College, Oberlin, OH, USA. ²Department of Psychology, Villanova University, Villanova, PA, USA. Email: psimen@oberlin.edu

tertone duration was much shorter or much longer than the criterion—i.e., when the duration could be easily discriminated from the intermediate, criterion duration.

Reward prediction errors are only prediction errors to the extent that they are surprising, however. An animal's surprise about the arrival time of the second tone ought, therefore, to modulate the reward prediction error, and it did. The intertone interval was picked from a set of six durations, so that a mouse's surprise about the second tone should become smaller as the intertone interval becomes longer—a longer wait for the second tone makes its imminent arrival more likely. The animal's surprise ought therefore to decrease to 0 at the maximum intertone duration. Soares *et al.*



The dopamine clock hypothesis of how we sense the passing of time is challenged by the findings of Soares *et al.*

demonstrate that dopamine bursts locked to the second tone were modulated by both reward probability and temporal surprise. Further, they found ramping dopaminergic neuron activity during the intertone interval that declined in the same way as the surprise function.

A faster subjective sense of time should produce a more rapidly declining surprise function, by definition. However, this seems to conflict with the dopamine clock hypothesis: Faster clocks should lead to more rapid declines in dopamine, but declines in dopamine should slow down the clock. This apparent inconsistency may rest on a lack of knowledge about how (6–8) or where (9–11) time is represented in the brain. Does a subjective time estimate computed outside the SNc drive the surprise function (12)? Or does the midbrain dopamine surprise signal drive a timer outside the SNc, as the dopamine clock hypothesis suggests? Soares *et al.* observed that lower intertone dopaminergic neuron activity correlated with a faster clock, not a slower clock, in contrast to the predictions of the

dopamine-timing hypothesis. Purely observational methods cannot establish causality, so the authors also used optogenetic methods that allow precise timing of perturbations to manipulate dopamine neuron activity on a subset of timing trials. Their results were unambiguous: Dopaminergic neuron activity did indeed appear to modulate internal clock speed, supporting a causal role for dopamine in regulating the subjective time sense. However, rather than speeding up the internal clock, increased dopaminergic neuron activity appeared to slow it down.

How can these results be reconciled with the large body of data on dopamine's role in subjective time estimation? One possibility is that, in contrast to the spatially and cell type-specific targeting of Soares *et al.*, systemic dopamine manipulation alters cortical timing processes (13) in a different direction and to a greater degree than the SNc-striatum effects.

Although phasic dopamine release seems to fit the reinforcement learning story, tonic dopamine release seems to do something rather different than encoding a reward prediction error. Instead, it seems to control overall response vigor or motivation (14). These findings can be accommodated within a theory in which tonic and phasic dopamine represent different kinds of information. Yet findings of “quasi-tonic” ramping signals within individual trials of nontiming tasks (15) pose a challenge even to this compromise theory. Soares *et al.* similarly see “quasi-tonic” surprise signals that challenge the dopamine clock hypothesis. These results suggest that upward- and downward-ramping dopamine signals may prove essential to unifying the reinforcement learning and interval-timing literatures, resolving their internal inconsistencies, and clarifying dopamine's true role in shaping behavior. ■

REFERENCES

1. P.R. Montague *et al.*, *J. Neurosci.* **16**, 1936 (1996).
2. S. Soares *et al.*, *Science* **354**, 1273 (2016).
3. W.H. Meck, *J. Exp. Psychol. Anim. Behav. Process.* **9**, 171 (1983).
4. J.R. Hollerman, W. Schultz, *Nat. Neurosci.* **1**, 304 (1998).
5. J.I. Lake, W.H. Meck, *Neuropsychologia* **51**, 284 (2013).
6. R. Laje, D. V. Buonomano, *Nat. Neurosci.* **16**, 925 (2013).
7. M. Matell, W.H. Meck, *Cogn. Brain Res.* **21**, 139 (2004).
8. P. Simen *et al.*, *Timing Time Percept.* **1**, 159 (2013).
9. A.A. Chubykin *et al.*, *Neuron* **77**, 723 (2013).
10. M. Wiener *et al.*, *NeuroImage* **49**, 1728 (2010).
11. M. Jazayeri, M.N. Shadlen, *Curr. Biol.* **25**, 2599 (2015).
12. Y.K. Takahashi *et al.*, *Neuron* **91**, 182 (2016).
13. K.L. Parker *et al.*, *J. Neurosci.* **34**, 16774 (2014).
14. Y. Niv *et al.*, *Psychopharmacology (Berl.)* **191**, 507 (2007).
15. M.W. Howe *et al.*, *Nature* **500**, 575 (2013).

10.1126/science.aal4021

EVOLUTION

Swimming in polluted waters

Genomic data provide insights into the molecular basis for pollution tolerance of Atlantic killifish

By Michael Tobler and Zachary W. Culumber

Human activities alter ecosystems around the planet, often rendering environmental conditions unfavorable for plant and animal survival. In the salt marshes along North America's Atlantic coast, the influx of industrial waste has caused chemical pollutants to accumulate at lethal levels, causing the disappearance of many species from affected sites. Yet, multiple populations of the Atlantic killifish (*Fundulus heteroclitus*) have adapted to cope with levels of pollution orders of magnitude higher than those that members of the same species from unaffected habitats can tolerate. On page 1305 of this issue, Reid *et al.* (1) provide strong evidence that adaptation has occurred rapidly and through similar genetic changes in multiple populations of killifish that have independently colonized polluted habitats.

Large population sizes and high levels of genetic diversity help to explain why killifish can rapidly adapt to anthropogenic pollution. Standing genetic variation facilitated rapid adaptation to toxic environments by recruiting existing, beneficial genetic variants, avoiding potential time lags that could occur if evolutionary responses were dependent on de novo beneficial mutations (2). Consequently, abundant species with high levels of genetic diversity may have the greatest ability to adapt to anthropogenic environmental change, whereas those lacking a standing stock of potentially beneficial mutations may be at risk of disappearing before evolution has a chance of rescuing them through adaptation.

In their study, Reid *et al.* compared whole-genome sequences from multiple pairs of tolerant and nontolerant populations. This

Division of Biology, Kansas State University, 116 Ackert Hall, Manhattan, KS 66506, USA. Email: tobler@ksu.edu

Why does time seem to fly when we're having fun?

Patrick Simen and Matthew Matell

Science **354** (6317), 1231-1232.
DOI: 10.1126/science.aal4021

ARTICLE TOOLS

<http://science.sciencemag.org/content/354/6317/1231>

RELATED CONTENT

<http://science.sciencemag.org/content/sci/354/6317/1273.full>

REFERENCES

This article cites 15 articles, 3 of which you can access for free
<http://science.sciencemag.org/content/354/6317/1231#BIBL>

PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)

Science (print ISSN 0036-8075; online ISSN 1095-9203) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. 2017 © The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. The title *Science* is a registered trademark of AAAS.