

Seroevidence for H5N1 Influenza Infections in Humans: Meta-Analysis

Taia T. Wang,¹ Michael K. Parides,² Peter Palese^{1,3*}

¹Department of Microbiology, Mount Sinai School of Medicine, New York, NY 10029, USA. ²Mount Sinai Center for Biostatistics and Department of Health Evidence and Policy, Mount Sinai School of Medicine, New York, NY 10029, USA.

³Department of Medicine, Mount Sinai School of Medicine, New York, NY 10029, USA.

*To whom correspondence should be addressed. E-mail: peter.palese@mssm.edu

Since 2003, there have been 573 World Health Organization (WHO) documented cases of avian H5N1 influenza infections in humans reported from 15 countries (1). Of the WHO confirmed cases, 58.6% have resulted in death (as of December 15th, 2011) (1, 2). These severe H5N1 infections were diagnosed using criteria developed by the WHO that are specific for H5N1 disease, but that lack the sensitivity to identify the total number of human infections (table S1) (3).

Given the fact that most H5N1 infections in poultry and in humans occur in resource-poor areas where access to health care is often arduous and expensive to obtain, we hypothesized that many people with H5N1 virus infection would not have been examined by a health provider to allow formal H5N1 disease confirmation. In addition, persons who are seropositive for H5N1 infection often report no history of influenza-like illness, and subclinical or mild H5N1 infections are not recognized under the WHO criteria for confirmed cases (4–8).

We conducted a meta-analysis of studies that evaluated the serological evidence of H5N1 infections in humans (table S2). The study participants, by and large, reported no recent respiratory and/or febrile illness. Variation between studies was taken into account using a random effects approach (9). We included in our primary analysis only studies that assessed serum samples based on modified WHO guidelines ($N=19$ study groups), or that presented data in such a way that modified WHO guidelines for H5N1 seropositivity could be applied [for modified WHO guidelines, see (10)]. In a secondary analysis, we compiled data from the remaining manuscripts based on the authors' criteria for positivity ($N=10$). Study participants with confirmed H5N1 infection were excluded to allow an analysis of seroprevalence in the persons without WHO-documented infection; exposure to poultry or humans with confirmed or suspected H5N1 infection was not a criterion for exclusion.

The primary analysis using WHO criteria was based on 7,304 study participants. All studies reported rates of seropositivity ranging from 0-5.3% with the exception of one study reporting 11.7% positivity among household contacts of infected individuals (5). Using WHO criteria, meta-analysis

revealed an overall seropositivity rate of 1.2% with a 95% confidence interval of 0.6% - 2.1% (Fig. 1A). Analysis of studies, which could not be interpreted by WHO guidelines [(10), table S2, and references therein], included 6,774 participants and yielded a seropositivity rate of 1.9% with a 95% confidence interval of 0.5 - 3.4% (Fig. 1B). Using either criterion, the rate of human H5N1 infections within the study populations was approximately 1-2% (10).

Using WHO criteria, we performed sub-analyses of study participants who were specifically employed as poultry workers ($N=2,729$) (4) (table S2). This analysis revealed a seropositivity rate of approximately 1.4%. If reports from the 1997 outbreak in Hong Kong are considered separately, the rate of seropositivity is approximately 3.2% (4, 5, 7). Studies after 1997, and that use WHO criteria, show an overall seropositivity rate of approximately 0.5% (8) (table S2).

The data were compiled from 12,677 study participants in 20 studies. They show that avian H5N1 viruses can cause a rate of mild or subclinical infections in humans that is not currently accounted for and thus, the true fatality rate for H5N1 influenza viruses is likely to be less than the frequently reported rate of more than 50%. Although it is not possible to determine an accurate fatality rate for H5N1 infections based on the data presented here, if one assumes a 1-2% infection rate in exposed populations, this would likely translate into millions of people who have been infected, worldwide. It is possible that deaths caused by H5N1 infection, as documented by the WHO, are also underestimated. We suggest that further investigation, on a large scale and by a standardized approach, is warranted to better estimate the total number of H5N1 infections that have occurred in humans. This information is critical for calculation of a real fatality rate that is not solely based on hospitalized patients.

References and Notes

1. WHO, Cumulative number of confirmed human cases for avian influenza A(H5N1) reported to WHO, 2003-2011; http://www.who.int/influenza/human_animal_interface/H5N1_cumulative_table_archives/en/index.html. (2011).

2. A. N. Abdel-Ghafar *et al.*, Update on avian influenza A (H5N1) virus infection in humans. *N Engl J Med* **358**, 261 (2008).
3. WHO, WHO case definitions for human infections with influenza A(H5N1) virus. (2006).
4. C. B. Bridges *et al.*, Risk of influenza A (H5N1) infection among poultry workers, Hong Kong, 1997, and 1998. *Journal of Infectious Diseases* **185**, 1005 (2002).
5. J. M. Katz *et al.*, Antibody response in individuals infected with avian influenza A (H5N1) viruses and detection of anti-H5 antibody among household and social contacts. *J Infect Dis* **180**, 1763 (1999).
6. P. Buchy *et al.*, Kinetics of neutralizing antibodies in patients naturally infected by H5N1 virus. *PLoS One* **5**, e10864 (2010).
7. B. C. Buxton *et al.*, Risk of influenza A (H5N1) infection among health care workers exposed to patients with influenza A (H5N1), Hong Kong. *J Infect Dis* **181**, 344 (2000).
8. P. Cavailler *et al.*, Seroprevalence of anti-H5 antibody in rural Cambodia, 2007. *Journal of Clinical Virology* **48**, 123 (2010).
9. R. DerSimonian, N. Laird, Meta-analysis in clinical trials. *Controlled clinical trials* **7**, 177 (1986).
10. Materials and Methods in supplementary online-material.
11. W. Cai *et al.*, Protective measures and H5N1-seroprevalence among personnel tasked with bird collection during an outbreak of avian influenza A/H5N1 in wild birds, Ruegen, Germany, 2006. *BMC Infect. Dis.* **9**, 170 (2009).
12. M. Ceyhan *et al.*, Serosurveillance study on transmission of H5N1 virus during a 2006 avian influenza epidemic. *Epidemiol. Infect.* **138**, 1274 (2010).
13. R. Dejphichai *et al.*, Seroprevalence of antibodies to avian influenza virus A (H5N1) among residents of villages with human cases, Thailand, 2005. *Emerg. Infect. Dis.* **15**, 756 (2009).
14. B. P. Khuntirat *et al.*, Evidence for subclinical avian influenza virus infections among rural Thai villagers. *Clin. Infect. Dis.* **53**, e107 (2011).
15. C. Y. Lu *et al.*, Potential infections of H5N1 and H9N2 avian influenza do exist in Guangdong populations of China. *Chin. Med. J. (Engl.)* **121**, 2050 (2008).
16. M. Robert, R. B. Holle, V. Setiawaty, K. N. Pangesti, E. R. Sedyaningsih, Seroprevalence of avian influenza A/H5N1 among poultry farmers in rural Indonesia, 2007. *Southeast Asian J. Trop. Med. Public Health* **41**, 1095 (2010).
17. C. Schultsz *et al.*, Prevalence of antibodies against avian influenza A (H5N1) virus among cullers and poultry workers in Ho Chi Minh City, 2005. *PLoS ONE* **4**, e7948 (2009).
18. M. Wang, C. X. Fu, B. J. Zheng, Antibodies against H5 and H9 avian influenza among poultry workers in China. *N. Engl. J. Med.* **360**, 2583 (2009).
19. R. Zhang, X. Rong, W. Pan, T. Peng, Determination of serum neutralization antibodies against seasonal influenza A strain H3N2 and the emerging strains 2009 H1N1 and avian H5N1. *Scand. J. Infect. Dis.* **43**, 216 (2011).
20. S. Hinjoy *et al.*, Low frequency of infection with avian influenza virus (H5N1) among poultry farmers, Thailand, 2004. *Emerg. Infect. Dis.* **14**, 499 (2008).
21. J. R. Ortiz *et al.*, Lack of evidence of avian-to-human transmission of avian influenza A (H5N1) virus among poultry workers, Kano, Nigeria, 2006. *J. Infect. Dis.* **196**, 1685 (2007).
22. K. Santhia *et al.*, Avian influenza A H5N1 infections in Bali Province, Indonesia: a behavioral, virological and seroepidemiological study. *Influenza Other Respir. Viruses* **3**, 81 (2009).
23. A. Apisarnthanarak *et al.*, Seroprevalence of anti-H5 antibody among Thai health care workers after exposure to avian influenza (H5N1) in a tertiary care center. *Clin. Infect. Dis.* **40**, e16 (2005).
24. S. Vong *et al.*, Low frequency of poultry-to-human H5N1 virus transmission, southern Cambodia, 2005. *Emerg. Infect. Dis.* **12**, 1542 (2006).
25. S. Vong *et al.*, Risk factors associated with subclinical human infection with avian influenza A (H5N1) virus—Cambodia, 2006. *J. Infect. Dis.* **199**, 1744 (2009).
26. H. Wang *et al.*, Probable limited person-to-person transmission of highly pathogenic avian influenza A (H5N1) virus in China. *Lancet* **371**, 1427 (2008).
27. J. H. Beigel *et al.*, Avian influenza A (H5N1) infection in humans. *N. Engl. J. Med.* **353**, 1374 (2005).
28. T. Rowe *et al.*, Detection of antibody to avian influenza A (H5N1) virus in human serum by using a combination of serologic assays. *J. Clin. Microbiol.* **37**, 937 (1999).
29. R. Kitphati *et al.*, Kinetics and longevity of antibody response to influenza A H5N1 virus infection in humans. *Clin. Vaccine Immunol.* **16**, 978 (2009).
30. L. V. Hedges, I. Olkin, M. Statistiker, *Statistical methods for meta-analysis*. (Academic Press Orlando, FL, 1985).
31. C. Clopper, E. S. Pearson, The Use of Confidence or Fiducial Limits Illustrated in the Case of the Binomial. *Biometrika* **26**, 404 (1934).
32. I. Belmaker *et al.*, Adherence with oseltamivir chemoprophylaxis among workers exposed to poultry during avian influenza outbreaks in southern Israel. *Int. J. Infect. Dis.* **13**, 261 (2009).
33. T. Nguyen *et al.*, Lack of H5N1 avian influenza transmission to hospital employees, Hanoi, 2004. (2005).
34. S. Velumani *et al.*, A novel peptide ELISA for universal detection of antibodies to human H5N1 influenza viruses. *PLoS ONE* **6**, e20737 (2011).
35. M. Prabakaran *et al.*, Development of epitope-blocking ELISA for universal detection of antibodies to human H5N1 influenza viruses. *PLoS ONE* **4**, e4566 (2009).
36. K. F. Shortridge, Pandemic influenza: a zoonosis? *Semin. Respir. Infect.* **7**, 11 (1992).

37. J. R. Ortiz *et al.*, No evidence of avian influenza A (H5N1) among returning US travelers. *Emerg. Infect. Dis.* **13**, 294 (2007).
38. N. T. Liem, W. Lim, Lack of H5N1 avian influenza transmission to hospital employees, Hanoi, 2004. *Emerg. Infect. Dis.* **11**, 210 (2005).
39. M. Wang *et al.*, Food markets with live birds as source of avian influenza. *Emerg. Infect. Dis.* **12**, 1773 (2006).
40. C. Schultsz *et al.*, Avian influenza H5N1 and healthcare workers. *Emerg. Infect. Dis.* **11**, 1158 (2005).

Acknowledgments: The authors thank Natalie Pica for her generous assistance in the preparation of this manuscript and Haomiao Jia for statistical analysis. This work was partially supported by National Institutes of Health (NIH) Grants U54 AI057158-04 and HHSN2662000700010C. Taia T. Wang was supported by NIH Training Grant T32 AI007647 and Mount Sinai Medical Scientists Training Grant T32 GM007280.

Supporting Online Material

www.sciencemag.org/cgi/content/full/science.1218888/DC1

Materials and Methods

Tables S1 and S2


References and Notes (11–40)

09 January 2012; accepted 15 February 2012


Published online 23 February 2012; 10.1126/science.1218888

Figure 1. Forest plots of overall, and study-specific, seroprevalence estimates with 95% confidence limits. Analysis using WHO criteria (A) or other criteria (see SOM) (B) for seropositivity. Individual values for prevalence (seropositivity), lower and upper confidence limits are shown at the right of each plot.

Estimates of Seropositivity (WHO Criteria)
Prevalence and 95% Confidence Limits



Estimates of Seropositivity (Other Criteria)
Prevalence and 95% Confidence Limits



Seroevidence for H5N1 Influenza Infections in Humans: Meta-Analysis

Taia T. Wang, Michael K. Parides and Peter Palese

published online February 23, 2012

ARTICLE TOOLS <http://science.sciencemag.org/content/early/2012/02/22/science.1218888>

SUPPLEMENTARY MATERIALS <http://science.sciencemag.org/content/suppl/2012/02/22/science.1218888.DC1>

RELATED CONTENT <http://science.sciencemag.org/content/sci/336/6088/1506.3.full>
<http://science.sciencemag.org/content/sci/336/6088/1506.2.full>

PERMISSIONS <http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)