

BIOTECHNOLOGY

CRISPR-Cas12a target binding unleashes indiscriminate single-stranded DNase activity


Janice S. Chen,^{1*} Enbo Ma,^{1*} Lucas B. Harrington,^{1*} Maria Da Costa,² Xinran Tian,³ Joel M. Palefsky,² Jennifer A. Doudna^{1,3,4,5,6,†}

CRISPR-Cas12a (Cpf1) proteins are RNA-guided enzymes that bind and cut DNA as components of bacterial adaptive immune systems. Like CRISPR-Cas9, Cas12a has been harnessed for genome editing on the basis of its ability to generate targeted, double-stranded DNA breaks. Here we show that RNA-guided DNA binding unleashes indiscriminate single-stranded DNA (ssDNA) cleavage activity by Cas12a that completely degrades ssDNA molecules. We find that target-activated, nonspecific single-stranded deoxyribonuclease (ssDNase) cleavage is also a property of other type V CRISPR-Cas12 enzymes. By combining Cas12a ssDNase activation with isothermal amplification, we create a method termed DNA endonuclease-targeted CRISPR trans reporter (DETECTR), which achieves attomolar sensitivity for DNA detection. DETECTR enables rapid and specific detection of human papillomavirus in patient samples, thereby providing a simple platform for molecular diagnostics.

CRISPR-Cas adaptive immunity in bacteria and archaea uses RNA-guided nucleases to target and degrade foreign nucleic acids (1–3). The CRISPR-Cas9 family of proteins has been widely deployed for gene-editing applications (4, 5) because of the precision of double-stranded DNA (dsDNA) cleavage induced by two catalytic domains, RuvC and HNH, at sequences complementary to a guide RNA (6, 7). A second family of enzymes, CRISPR-Cas12a (Cpf1), uses a single RuvC catalytic domain for guide RNA-directed dsDNA cleavage (8–13) (Fig. 1A). Distinct from Cas9, Cas12a enzymes recognize a T nucleotide-rich protospacer-adjacent motif (PAM) (8), catalyze their own guide CRISPR RNA (crRNA) maturation (14), and generate a PAM-distal dsDNA break with staggered 5' and 3' ends (8), features that have attracted interest for gene-editing applications (15, 16). However, the substrate specificity and DNA cleavage mechanism of Cas12a remain to be fully elucidated.

While investigating substrate requirements for Cas12a activation, we tested *Lachnospiraceae* bacterium ND2006 Cas12a (LbCas12a) for guide RNA-directed single-stranded DNA (ssDNA) cleavage, a capability of diverse CRISPR-Cas9 orthologs (17, 18). Purified LbCas12a or *Streptococcus pyogenes* Cas9 (SpCas9) proteins (fig. S1) were

assembled with guide RNA sequences targeting a circular, single-stranded M13 DNA phage. In contrast to SpCas9, we were surprised to find that LbCas12a induced rapid and complete degradation of M13 by a cleavage mechanism that could not be explained by sequence-specific DNA cutting (Fig. 1B). This ssDNA shredding activity, not observed with a catalytically inactive LbCas12a (D832A, with an Asp⁸³²→Ala substitution), raised



¹Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, CA 94720, USA. ²Department of Medicine, University of California, San Francisco, San Francisco, CA 94143, USA. ³Department of Chemistry, University of California, Berkeley, Berkeley, CA 94720, USA. ⁴Innovative Genomics Institute, University of California, Berkeley, Berkeley, CA 94704, USA. ⁵Howard Hughes Medical Institute, University of California, Berkeley, Berkeley, CA 94720, USA. ⁶Molecular Biophysics and Integrated Bioimaging Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA.

*These authors contributed equally to this work.

†Corresponding author. Email: doudna@berkeley.edu

the possibility that a target-bound LbCas12a could degrade any ssDNA sequence. Remarkably, LbCas12a also catalyzed M13 degradation in the presence of a different guide RNA and its complementary ssDNA “activator” that have no sequence homology to the M13 phage genome (Fig. 1C). These findings reveal that binding of the LbCas12a-crRNA complex to a guide-complementary ssDNA unleashes robust, nonspecific ssDNA trans-cleavage activity.

We next investigated the requirements for LbCas12a-catalyzed trans-cleavage activity. Using a fluorophore quencher (FQ)-labeled reporter assay (19), we assembled LbCas12a with its crRNA and either a complementary ssDNA, dsDNA, or single-stranded RNA (ssRNA) and introduced an unrelated ssDNA- or ssRNA-FQ reporter in trans (fig. S2). Both the crRNA-complementary ssDNA or dsDNA (the activator) triggered LbCas12a to cleave the ssDNA-FQ reporter substrate (fig. S2A). However, ssRNA was neither capable of activating trans cleavage nor susceptible to degradation by LbCas12a (fig. S2B), confirming that LbCas12a harbors DNA-activated general deoxyribonuclease (DNase) activity.

To determine how LbCas12a-catalyzed ssDNA cleavage relates to site-specific dsDNA cutting, we tested the target strand (TS) and nontarget strand (NTS) requirements for LbCas12a activation. Although TS cutting occurred irrespective of the NTS length (fig. S3A), NTS cleavage occurred only when the TS contained at least 15 nucleotides of complementarity with the crRNA (fig. S3B). This shows that TS recognition is a prerequisite for NTS cutting. To test whether LbCas12a remains active for nonspecific ssDNA cleavage after

dsDNA target cleavage, we first cut a dsDNA plasmid with LbCas12a-crRNA and then added an unrelated dsDNA or ssDNA to the reaction (Fig. 2A). Whereas the nonspecific dsDNA substrate remained intact, the ssDNA was rapidly degraded in a RuvC domain-dependent manner (Fig. 2A and figs. S4 and S5). Using truncated activators that are too short to be cleaved, we determined that only target-DNA binding is required to activate trans-ssDNA cleavage (fig. S6). Together, these results show that RNA-guided DNA binding activates LbCas12a for both site-specific dsDNA cutting and nonspecific ssDNA trans cleavage.

The rapid degradation of a trans substrate suggested that the kinetics of LbCas12a-catalyzed site-specific dsDNA (cis) cleavage and nonspecific ssDNA (trans) cleavage are fundamentally different. Stoichiometric titration experiments showed

that cis cleavage is single turnover (20) (Fig. 2B), whereas trans cleavage is multiple turnover (Fig. 2C). Although the LbCas12a-crRNA complex remains bound to the dsDNA target after cis cleavage, the complex releases its PAM-distal cleavage products from the RuvC active site (20), enabling ssDNA substrate access and turnover. We found that LbCas12a-crRNA bound to a ssDNA activator molecule catalyzed trans-ssDNA cleavage at a rate of ~250 turnovers per second and a catalytic efficiency (k_{cat}/K_m) of $5.1 \times 10^8 \text{ s}^{-1} \text{ M}^{-1}$. When bound to a dsDNA activator, LbCas12a-crRNA catalyzed ~1250 turnovers per second with a catalytic efficiency approaching the rate of diffusion (21), a k_{cat}/K_m of $1.7 \times 10^9 \text{ s}^{-1} \text{ M}^{-1}$ (Fig. 2D and fig. S7). These differences suggest that the NTS of the dsDNA activator helps stabilize the Cas12a complex in an optimal conformation for trans-ssDNA cutting.

We next tested the specificity of trans-cleavage activation using either a ssDNA or dsDNA activator. We found that the PAM sequence required for dsDNA binding by CRISPR-Cas12a (20) is critical for catalytic activation by a crRNA-complementary dsDNA (9) but not for a crRNA-complementary ssDNA (Fig. 3A). Mismatches between the crRNA and activator slowed the trans-cleavage rate, with PAM mutations or mismatches in the PAM-adjacent “seed region” inhibiting trans-ssDNA cleavage only for a dsDNA activator (Fig. 3B and fig. S8). Consistent with the mismatch tolerance pattern observed in Cas12a off-target studies (22, 23), these data corroborate PAM-mediated target-dsDNA binding and the role of base pairing between the crRNA and target strand to activate trans-ssDNA cutting.

We wondered if this trans-ssDNA-cutting activity might be a property shared by type V




Fig. 2. Kinetics of Cas12a ssDNA trans cleavage. (A) Sequence-specific plasmid DNA cleavage reactions by LbCas12a-crRNA complex (top) were introduced to a separate radiolabeled dsDNA or ssDNA substrate of unrelated sequence (bottom); time course represented in minutes. Substrate (S) and nucleotide products (P) were resolved by agarose gel electrophoresis (top) or denaturing polyacrylamide gel electrophoresis (PAGE) (bottom). (B) Target dsDNA or (C) nonspecific ssDNA incubated with molar ratios of LbCas12a-crRNA, as indicated. Each point represents

the mean quantified percent cleavage after 30 min at 37°C, at which time the reaction was at completion. Error bars represent mean \pm SD, where $n = 3$ replicates. Yellow coloring indicates PAM. Arrowheads indicate cleavage sites. (D) Representative Michaelis-Menten plot for LbCas12a-catalyzed ssDNA trans cleavage using a dsDNA or ssDNA activator. Measured k_{cat}/K_m values reported as mean \pm SD, where $n = 3$ Michaelis-Menten fits. V_0 , rate of catalysis. Color scheme in (B) and (C) is the same as for (D). In (C), blue indicates nonspecific DNA.





Fig. 3. Specificity and conservation of trans-cleavage activation. (A) LbCas12a-crRNA in the absence or presence of indicated activator, incubated with a radiolabeled nonspecific ssDNA substrate (S) for 30 min at 37°C; products (P) resolved by denaturing PAGE. (B) Observed trans-cleavage rates for LbCas12a using a ssDNA or dsDNA activator with indicated mismatches; rates represent the average of three different targets measured in triplicate, and error bars represent mean \pm SD, where

$n = 9$ (three replicates for three independent targets). PT, perfect target; mut PAM, mutated PAM. (C) Radiolabeled cis (complementary) or trans (noncomplementary) substrates were incubated with Cas12a-crRNA or Cas9-sgRNA in the presence or absence of a ssDNA activator for 30 min at 37°C; a cis-dsDNA substrate was used in the “no enzyme” lanes. Substrate (S) and nucleotide products (P) were resolved by denaturing PAGE. NmCas9 is *Neisseria meningitidis* Cas9.

Fig. 4. Rapid identification of HPV16 and HPV18 in human samples by DETECTR.

(A) Diagram of HPV16 and HPV18 sequences within the hypervariable loop V of the L1-encoding gene targeted by Cas12a; yellow highlighted bases indicate 5' PAM sequence. (B) Heatmap represents normalized mean fluorescence values of HPV16 and HPV18 detected in human cell lines by DETECTR; normalized scale shown in (D). (C) Schematic outlining DNA extraction from human anal samples to HPV identification by DETECTR. RPA, recombinase polymerase amplification; F, fluorophore; Q, quencher. (D) Identification of HPV16 and HPV18 in 25 patient samples by PCR (left) and DETECTR (right). DETECTR heatmap represents normalized mean fluorescence values.



CRISPR effector proteins, considering that all Cas12 proteins contain a single RuvC nuclease domain (3, 24). Consistent with this possibility, purified Cas12a orthologs from *Acidaminococcus* sp. (AsCas12a) and *Francisella novicida* (FnCas12a), as well as a Cas12b protein from *Alicyclobacillus acidoterrestris* (AaCas12b), all catalyzed non-specific ssDNA cleavage when assembled with a crRNA and complementary ssDNA activator (Fig. 3C and fig. S9). By contrast, none of the type II CRISPR-Cas9 proteins tested showed evidence for trans cleavage (Fig. 3C and fig. S9). These results reveal the unexpected functional convergence of Cas12 enzymes with the type III CRISPR-Csm/Cmr and type VI CRISPR-Cas13 effectors, which also exhibit target-activated, nonspecific ssDNase or single-stranded ribonuclease (ssRNase) activity, respectively (25, 26).

We next explored whether LbCas12a could be repurposed as a DNA-detection platform. In particular, accurate and rapid detection of human papillomavirus (HPV) is critical for identifying those at risk of HPV-related cancers, with types 16 (HPV16) and 18 (HPV18) accounting for most precancerous lesions (27). To test if LbCas12a could distinguish between these two dsDNA viruses, we selected a target sequence next to a TTTA PAM that varied by six base pairs between the two HPV genotypes (fig. S10). HPV16- or

HPV18-containing plasmids were incubated with LbCas12a-crRNA that targeted either the HPV16 or HPV18 fragment and a ssDNA-FQ reporter, which produced a signal only in the presence of the cognate target (fig. S10). To enhance sensitivity, we coupled isothermal amplification by recombinase polymerase amplification with LbCas12a to develop a one-pot detection method termed DNA endonuclease-targeted CRISPR trans reporter (DETECTR) (fig. S11A). When programmed to its cognate plasmid, DETECTR identified targets with attomolar sensitivity (fig. S11B).

To assess whether we could detect HPV in complex mixtures, DNA extracted from cultured human cells infected with HPV16 (SiHa cells) or HPV18 (HeLa cells) or without HPV (BJAB cells) was added to LbCas12a-crRNA targeting the hypervariable loop V of the L1-encoding gene within HPV16 or HPV18 (Fig. 4A). Whereas LbCas12a-crRNA alone lacked the sensitivity to detect HPV, DETECTR unambiguously identified HPV16 and HPV18 only in SiHa and HeLa cells, respectively (Fig. 4B and fig. S12, A and B). To investigate the utility of DETECTR for analyzing patient samples, we tested crude DNA extractions from 25 human anal swabs previously analyzed by a polymerase chain reaction (PCR)-based method for HPV infection (fig. S13) (28). Within 1 hour, DETECTR accurately identified HPV16

(25/25 agreement) and HPV18 (23/25 agreement) in patient samples containing a heterogeneous mixture of HPV types, with good correlation between the PCR-based intensity and DETECTR signal (Fig. 4, C and D, and figs. S12, C and D, and S13). These results demonstrate a new platform for CRISPR-based diagnostics, similar to that developed for RNA detection using CRISPR-Cas13a (19, 29), and suggest that DETECTR could, in principle, detect any DNA sequence with high sensitivity and specificity (fig. S14).

Together, these findings support a mechanism of target interference that begins with the Cas12a-guide RNA complex binding to a complementary DNA sequence in a PAM-dependent (dsDNA) or PAM-independent (ssDNA) manner (fig. S15). Within a host bacterium, such enzyme activation could provide simultaneous protection from both dsDNA and ssDNA phages and could also target ssDNA sequences that arise temporarily during phage replication or transcription (30). In a genome-editing context, target-activated ssDNA cutting by Cas12a has the potential to cleave transiently exposed ssDNA at replication forks (31), R-loops (32), and transcription bubbles (33) or ssDNA templates used for homology-directed repair (34). Finally, unleashing the ssDNase activity of Cas12 proteins offers a new strategy to improve the speed, sensitivity, and specificity of molecular diagnostic applications.

REFERENCES AND NOTES

- R. Barrangou *et al.*, *Science* **315**, 1709–1712 (2007).
- L. A. Marraffini, E. J. Sontheimer, *Science* **322**, 1843–1845 (2008).
- E. V. Koonin, K. S. Makarova, F. Zhang, *Curr. Opin. Microbiol.* **37**, 67–78 (2017).
- J. A. Doudna, E. Charpentier, *Science* **346**, 1258096 (2014).
- R. Barrangou, P. Horvath, *Nat. Microbiol.* **2**, 17092 (2017).
- M. Jinek *et al.*, *Science* **337**, 816–821 (2012).
- J. S. Chen, J. A. Doudna, *Nat. Rev. Chem.* **1**, 0078 (2017).
- B. Zetsche *et al.*, *Cell* **163**, 759–771 (2015).
- D. C. Swarts, J. van der Oost, M. Jinek, *Mol. Cell* **66**, 221–233, e4 (2017).
- D. Dong *et al.*, *Nature* **532**, 522–526 (2016).
- P. Gao, H. Yang, K. R. Rajashankar, Z. Huang, D. J. Patel, *Cell Res.* **26**, 901–913 (2016).
- S. Stella, P. Alcón, G. Montoya, *Nature* **546**, 559–563 (2017).
- T. Yamano *et al.*, *Cell* **165**, 949–962 (2016).
- I. Fonfara, H. Richter, M. Bratovič, A. Le Rhun, E. Charpentier, *Nature* **532**, 517–521 (2016).
- B. Zetsche *et al.*, *Nat. Biotechnol.* **35**, 31–34 (2017).
- X. Tang *et al.*, *Nat. Plants* **3**, 17018 (2017).
- Y. Zhang, R. Rajan, H. S. Seifert, A. Mondragón, E. J. Sontheimer, *Mol. Cell* **60**, 242–255 (2015).
- E. Ma, L. B. Harrington, M. R. O’Connell, K. Zhou, J. A. Doudna, *Mol. Cell* **60**, 398–407 (2015).
- A. East-Seletsky *et al.*, *Nature* **538**, 270–273 (2016).
- D. Singh, J. Mallon, A. Poddar, Y. Wang, R. Tipanna, O. Yang, S. Bailey, T. Ha, Real-time observation of DNA target interrogation and product release by the RNA-guided endonuclease CRISPR Cpf1. *BioRxiv* 205575 [Preprint], 18 October 2017.
- R. A. Alberty, G. G. Hammes, *J. Phys. Chem.* **62**, 154–159 (1958).
- D. Kim *et al.*, *Nat. Biotechnol.* **34**, 863–868 (2016).
- B. P. Kleinstiver *et al.*, *Nat. Biotechnol.* **34**, 869–874 (2016).
- S. Shmakov *et al.*, *Nat. Rev. Microbiol.* **15**, 169–182 (2017).
- J. R. Elmore *et al.*, *Genes Dev.* **30**, 447–459 (2016).
- O. O. Abudayyeh *et al.*, *Science* **353**, aaf5573 (2016).
- C. B. Woodman, S. I. Collins, L. S. Young, *Nat. Rev. Cancer* **7**, 11–22 (2007).

28. J. M. Palefsky, E. A. Holly, M. L. Ralston, N. Jay, *J. Infect. Dis.* **177**, 361–367 (1998).
29. J. S. Gootenberg *et al.*, *Science* **356**, 438–442 (2017).
30. P. Samai *et al.*, *Cell* **161**, 1164–1174 (2015).
31. J. M. Sogo, M. Lopes, M. Foiani, *Science* **297**, 599–602 (2002).
32. V. Bhatia, E. Herrera-Moyano, A. Aguilera, B. Gómez-González, *Genes (Basel)* **8**, E171 (2017).
33. Y. Zhang *et al.*, *Science* **338**, 1076–1080 (2012).
34. C. D. Richardson, G. J. Ray, M. A. DeWitt, G. L. Curie, J. E. Corn, *Nat. Biotechnol.* **34**, 339–344 (2016).

ACKNOWLEDGMENTS

We thank O. Mavrothalassitis, D. Burstein, D. Pérez-Espino, G. Knott, A. Wright, J. Cofsky, D. Lee, and members of the Doudna laboratory for comments and discussions, and we thank the patients for their generosity. **Funding:** This research was supported in part by the Allen Distinguished Investigator Program through The Paul G. Allen Frontiers Group and the NSF (MCB-1244557 to J.A.D.). J.S.C. and L.B.H. are

supported by NSF Graduate Research Fellowships. **Author contributions:** J.S.C., E.M., and L.B.H. contributed equally to the work and conceived of and designed experiments. J.S.C., E.M., and L.B.H. performed protein expression and biochemical experiments, with assistance from X.T. J.S.C. designed and performed DETECTR assays, with input from M.D.C. and J.M.P. regarding HPV detection in human cell lines and patient samples. J.S.C., E.M., L.B.H., and J.A.D. wrote the manuscript. **Competing interests:** J.S.C. and L.B.H. are cofounders of Mammoth Biosciences. J.M.P. is a professor of medicine at the University of California, San Francisco, and a scientific adviser to Antiva Biosciences, Agenovir, and Merck and Co. J.A.D. is an investigator of the Howard Hughes Medical Institute and executive director of the Innovative Genomics Institute at the University of California, Berkeley, and the University of California, San Francisco. J.A.D. is a cofounder of Editas Medicine, Intellia Therapeutics, Scribe Therapeutics, and Caribou Biosciences; a scientific adviser to Caribou, Intellia,

eFFECTOR Therapeutics, Scribe Therapeutics, Synthego, Metagenomi, Mammoth Biosciences, and Driver; and is on the Board of Directors of Driver and Johnson & Johnson. The Regents of the University of California have patents pending for CRISPR technologies on which the authors are inventors. **Data and materials availability:** All data are available in the manuscript or the supplementary materials.

SUPPLEMENTARY MATERIALS

www.sciencemag.org/content/360/6387/436/suppl/DC1
Materials and Methods
Figs. S1 to S15
Table S1
References (35–37)

29 November 2017; resubmitted 22 January 2018
Accepted 5 February 2018
Published online 15 February 2018
10.1126/science.aar6245

CRISPR-Cas12a target binding unleashes indiscriminate single-stranded DNase activity

Janice S. Chen, Enbo Ma, Lucas B. Harrington, Maria Da Costa, Xinran Tian, Joel M. Palefsky and Jennifer A. Doudna

Science **360** (6387), 436-439.

DOI: 10.1126/science.aar6245 originally published online February 15, 2018

Taking CRISPR technology further

CRISPR techniques are allowing the development of technologies for nucleic acid detection (see the Perspective by Chertow). Taking advantages of the distinctive enzymatic properties of CRISPR enzymes, Gootenberg *et al.* developed an improved nucleic acid detection technology for multiplexed quantitative and highly sensitive detection, combined with lateral flow for visual readout. Myhrvold *et al.* added a sample preparation protocol to create a field-deployable viral diagnostic platform for rapid detection of specific strains of pathogens in clinical samples. Cas12a (also known as Cpf1), a type V CRISPR protein, cleaves double-stranded DNA and has been adapted for genome editing. Chen *et al.* discovered that Cas12a also processes single-stranded DNA threading activity. A technology platform based on this activity detected human papillomavirus in patient samples with high sensitivity.

Science, this issue p. 439, p. 444, p. 436; see also p. 381

ARTICLE TOOLS

<http://science.sciencemag.org/content/360/6387/436>

SUPPLEMENTARY MATERIALS

<http://science.sciencemag.org/content/suppl/2018/02/14/science.aar6245.DC1>

RELATED CONTENT

<http://science.sciencemag.org/content/sci/360/6387/439.full>
<http://science.sciencemag.org/content/sci/360/6387/381.full>
<http://science.sciencemag.org/content/sci/360/6387/444.full>
<http://stm.sciencemag.org/content/scitransmed/9/372/eaah3480.full>
<http://stm.sciencemag.org/content/scitransmed/9/418/eaan8081.full>
<http://stm.sciencemag.org/content/scitransmed/8/360/360ra134.full>
file:/content

REFERENCES

This article cites 36 articles, 11 of which you can access for free
<http://science.sciencemag.org/content/360/6387/436#BIBL>

PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)