

tissues was greatly decreased in pancreatic diabetes and accordingly suggested that the defective oxidation in this type of diabetes was due to the decrease in catalase. If the defective oxidation in diabetes is due to the decrease in the catalase of the tissues and if it can be shown that the administration of alcohol produces an increase in the catalase of the tissues due to the stimulation of the liver to an increased output of this enzyme into the blood, then it would seem probable that the helpful effect of alcohol in diabetes is due to the increase in catalase with resulting increase in oxidation.

Dogs were used in the investigation. The catalase in 0.5 c.c. of the blood of the animals was determined by adding this amount of blood to 50 c.c. of hydrogen peroxide in a bottle at 22° C. and as the oxygen gas was liberated, it was conducted through a rubber tube to an inverted burette previously filled with water. After the volume of gas thus collected in ten minutes had been reduced to standard atmospheric pressure, the resulting volume was taken as a measure of the amount of catalase in the 0.5 c.c. of blood. The material was shaken at a fixed rate of one hundred and eighty double shakes per minute during the determinations.

Twenty-five c.c. per kilo of body weight of 45 per cent. ethyl alcohol were introduced into the stomachs of the animals by means of a stomach tube. Previous to as well as at 15-minute intervals after the introduction of alcohol, the catalase in 0.5 c.c. of blood taken from the external jugular was determined. Fifteen minutes after the introduction of alcohol into the stomach, it was found that the catalase of the blood was increased by about 30 per cent., after 30 minutes by about 50 per cent., and after 45 minutes the catalase of the blood of some of the dogs was increased by as much as 100 per cent.

After etherizing other dogs the abdominal wall was opened and the catalase of the blood taken directly from the liver or from one of the hepatic veins as well as from the jugular was determined. It was found that the blood from the liver was richer in catalase by ten to

fifteen per cent. than the blood from any other part of the body. This was taken to mean that there is a continuous output of catalase from the liver into the blood and that this catalase is taken to the tissues to be used presumably in the oxidative processes. After the introduction of the alcohol into the stomach of the animal, it was found that the catalase in the blood taken directly from the liver was increased much more rapidly than that taken from a systemic vein such as the jugular, hence the alcohol must have been stimulating the liver to an increased output of catalase and in this manner producing an increase in the catalase of the blood and hence of the tissues.

Alcohol was also administered to dogs rendered diabetic by the removal of the pancreas, and it was found that the catalase of the blood and hence of the tissues of these animals was increased. It is probable that the helpful effect of alcohol in states of depression and in convalescence as well as the exhilarating effect on normal subjects is due to the stimulation of the liver to an increased output of catalase with resulting increase in oxidation.

The conclusion is drawn that the administration of alcohol to diabetics is helpful because it stimulates the liver to an increased output of catalase which is carried by the blood to the tissues where it facilitates the oxidative processes with resulting increased oxidation of sugar and decreased acidosis.

W. E. BURGE

PHYSIOLOGICAL LABORATORY OF THE
UNIVERSITY OF ILLINOIS

SCIENCE

A Weekly Journal devoted to the Advancement of
Science, publishing the official notices and pro-
ceedings of the American Association for
the Advancement of Science

Published every Friday by

THE SCIENCE PRESS
LANCASTER, PA. GARRISON, N. Y.
NEW YORK, N. Y.

Entered in the post-office at Lancaster, Pa., as second class matter