

SCIENCE

FRIDAY, APRIL 15, 1921

CONTENTS

<i>The Photochemistry of the Sensitivity of Animals to Light:</i> DR. SELIG HECHT	347
<i>The Mechanism of Injury and Recovery of the Cell:</i> PROFESSOR W. J. V. OSTERHOUT.	356
Isao Iijima: PROFESSOR BASHFORD DEAN	356
<i>Scientific Events:—</i>	
<i>Ex-secretary Meredith on Research; Scientific Legislation; Lectures before the Sioux City Academy; Cooperation of National Health Agencies; The American Meteorological Society; The Edinburgh Meeting of the British Association</i>	357
<i>Scientific Notes and News</i>	360
<i>University and Educational News</i>	363
<i>Discussion and Correspondence:—</i>	
<i>Positive Ray Analysis of Lithium:</i> DR. A. J. DEMPSTER. <i>A Remedy for Mange in White Rats:</i> CORNELIA KENNEDY. <i>Impossible (?) Stories:</i> PROFESSOR SAMUEL G. BARTON	363
<i>Quotations:—</i>	
<i>International Scientific Organization</i>	364
<i>Scientific Books:—</i>	
<i>The Terrestrial Life associated with the Coals of Northern France:</i> PROFESSOR CHARLES SCHUCHERT	367
<i>Special Articles:—</i>	
<i>The Relativity Shift of Spectrum Lines:</i> DR. RAYMOND T. BIRGE. <i>A New High Temperature Record for Growth:</i> DR. D. T. MACDOUGAL	368
<i>The American Mathematical Society:</i> PROFESSOR R. G. D. RICHARDSON	372

THE PHOTOCHEMISTRY OF THE SENSITIVITY OF ANIMALS TO LIGHT¹

I

AN analysis of sensory stimulation, in order to be objective, must take its data from the relations between the properties of the stimulating agent and those of the responses of the animal. If the analysis is to be quantitative as well as objective, not only should the response be a qualitatively invariable reflex but, together with the source of stimulation, it should be capable of precise and easy control.

There are a number of animals which possess such characteristic responses. Typical of these are the ascidian, *Ciona intestinalis* and the long-neck clam, *Mya arenaria*. Both of these animals, when exposed to light, respond by a vigorous retraction of the siphons. It has therefore been possible to investigate quantitatively the properties of their photic sensitivity, and as a result to propose an hypothesis which accounts for this type of irritability in terms of an underlying photochemical mechanism.

I propose now to describe briefly the evidence which has been accumulated in this connection, and to present the outstanding features of the proposed hypothetical mechanism.

II

The photosensory activities of these animals possess four striking and important properties. (1) When exposed to light, the animal

¹Delivered at the Symposium on General Physiology held by the American Society of Naturalists on December 30, 1920, at its Chicago meetings. The paper was illustrated with a number of charts which are not reproduced here. They may be found, together with the data on which this summary is based, in a series of articles in the *Journal of General Physiology* from 1918 to the present time.

MSS. intended for publication and books, etc., intended for review should be sent to The Editor of Science, Garrison-on-Hudson, N. Y.

Science

53 (1372)

Science **53** (1372), 347-372.

ARTICLE TOOLS

<http://science.sciencemag.org/content/53/1372.citation>

PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)

Science (print ISSN 0036-8075; online ISSN 1095-9203) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. The title *Science* is a registered trademark of AAAS.

Copyright © 1921 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works.