

SCIENCE

FRIDAY, JULY 22, 1921

TYPES OF VALENCE

CONTENTS

<i>Types of Valence:</i> DR. IRVING LANGMUIR.....	59
<i>Professor H. Bruchmann:</i> PROFESSOR DOUGLAS HOUGHTON CAMPBELL	67
<i>Scientific Events:—</i>	
<i>Field Work of the Smithsonian Institution; The Medical School of the University of Virginia; The Science Club of the University of Texas; The Rochester Meeting of the Optical Society of America; American Engineers in Europe</i>	68
<i>Scientific Notes and News</i>	71
<i>University and Educational News</i>	72
<i>Discussion and Correspondence:—</i>	
<i>The Geographic Distribution of Hybrids: DR. M. L. FERNALD. Albinism in the Black Bear: PAUL C. STANDLEY. Bechhold's "Capillary Phenomenon" in Agriculture: JEROME ALEXANDER</i>	73
<i>Quotations:—</i>	
<i>The Royal Institution.....</i>	74
<i>Scientific Books:—</i>	
<i>Evermann and Clark on Lake Mazinkuckee: T. L. HANKINSON.....</i>	75
<i>Special Articles:—</i>	
<i>An Explanation of Liesegang's Rings: DR. HUGH MCGUIGAN</i>	78
<i>The North Carolina Academy of Science: C. S. BRIMLEY</i>	80

FROM the theory proposed by G. N. Lewis in 1916 and subsequently extended by the writer, it is clear¹ that the term valence has been used in the past to cover what we may now recognize as three distinct types of valence, viz.:

1. Positive valence: the number of electrons an atom can give up.
2. Negative valence: the number of electrons an atom can take up.
3. Covalence: the number of pairs of electrons an atom can share with its neighbors.

It was shown that these fundamental conceptions of valence as well as the actual numerical values of each type of valence for most of the elements could be derived from a few postulates regarding the structure of atoms. The following method of deriving these relationships, however, is not only much simpler than that previously given by the writer, but throws a new light on the relationships between the different types of valence.

We will take for granted the Rutherford type of atom, which consists of a positive nucleus surrounded by a number of electrons equal to the atomic number of the atom. We will also assume that Coulomb's law applies to the forces between the charged particles in the atom, but at the same time will recognize the existence of repulsive forces in atoms which prevent the electrons from falling into the nucleus. For the present purpose, however, it is immaterial whether the repulsive force is a dynamic force (centrifugal force) such as that assumed by Bohr, or is a static force as postulated by G. N. Lewis, J. J. Thomson or recently by the writer.²

We shall need to make only 3 postulates in

¹ Langmuir, *Jour. Amer. Chem. Soc.*, 41, 926 (1919), and *Jour. Ind. Eng. Chem.*, 12, 386 (1920).

² SCIENCE, 53, 290, Mar. 25, 1921.

Science

54 (1386)

Science **54** (1386), 59-80.

ARTICLE TOOLS

<http://science.sciencemag.org/content/54/1386.citation>

PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)

Science (print ISSN 0036-8075; online ISSN 1095-9203) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. The title *Science* is a registered trademark of AAAS.

Copyright © 1921 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works.