

is found to pass through a maximum as the distance of the ring from the x-axis increases. Our theory identifies the weight of this maximum volume below the surfaces with the maximum pull on the ring as found experimentally. Harkins, Young and Cheng plot a function of this maximum pull,  $R^3/\bar{V}$ , where  $R$  is the radius of the ring and  $V$  is the volume of the liquid equivalent to the weight measured by this pull, against  $\gamma/p$ , a curve for each value of  $\frac{R}{r}$ , where  $\gamma$  is the surface tension of the liquid as determined by a reference method, and  $p$  is the pull on the ring divided by  $4\pi R$ . We show that  $\gamma/p$  is equal to  $2\pi R/\bar{V}$ , a ratio which does not involve the determination of  $\gamma$  by any other method. Now the correction curves found by Harkins and his collaborators are identical with the curves calculated by us from the maximum volume below the surfaces and plotted as indicated, i.e.,  $R^3/\bar{V}$  by  $2\pi R/\bar{V}$  for a particular  $R/r$ . Our calculated curves are probably not as precise as the experimental ones because of the lack of sufficient calculated surfaces and the errors due to the consequent interpolations. But the excellent agreement that is obtained shows that the same values of  $\gamma$ , within the small error of both methods, are obtained whether the relative corrections of the Harkins procedure or the absolute corrections as calculated by us are used. Thus the ring method may now be considered an absolute one, since by it surface tension may be determined without reference to any other procedure. A detailed discussion of this theory is to be published soon.

B. B. FREUD  
H. Z. FREUD

#### PROLONGED EFFECT ON DIGITALIS PURPUREA OF EXPOSURE UNDER ULTRA-VIOLET-TRANSMITTING GLASS

For the past three years, a study has been made of the effect produced upon *Digitalis purpurea* L. by exposure in the greenhouse stage under an ultra-violet-transmitting glass. *D. purpurea* is well known as the common garden foxglove, and is also a very important medicinal plant. In 1927 and again in 1928, seedlings were grown about three months under an ultra-violet-transmitting glass, with controls under ordinary glass in an adjacent section of the same greenhouse. Thus it was possible to eliminate differences of temperature, moisture and other factors.

Both series were then grown in adjacent beds under open field culture conditions; and in each year, as reported<sup>1</sup> at the Nashville and New York meetings, the treated plants proved considerably superior to

their controls in the content of the characteristic active principles.

The final report now to be given concerns the history of the 1928 plants during their second season of growth, this species being biennial. By careful mulching they were brought through the winter in good condition. Our capricious Michigan weather in 1929 made it necessary to water the plants last summer throughout a season marked by severe drought; but all were treated the same, and had attained the normal early blooming stage at the time the samples were harvested.

For many years it was held that second-year leaves were more potent than first-year herbage but eventually this idea was refuted. We now know from numerous tests that greater activity occurs during the first year of a digitalis plant's life. Therefore, it is to be noted that the actual percentages now given, obtained from second-year growth, are not as high as those previously reported from first-year leaves; but the relative values are again confirmed; the treated plants have the higher potency. Exact values are as tabulated below, all methods having meticulously followed those heretofore used.

Treated group .....	250 per cent. of standard
Control group .....	165 per cent. of standard
	Increase of potency, 51.5 per cent.

It thus seems clearly established that *D. purpurea* is benefited permanently by exposure during seedling development under glass that affords an increased amount of ultra-violet radiation. Where digitalis passes its whole cycle under natural, out-of-door conditions in a mild climate, this fact may not be of great significance, but for successful culture of high-grade plants in Michigan we must depend in large measure upon indoor protection during the seedling stage. It is gratifying, therefore, to know that modern methods of glass making have increased the efficiency of greenhouse propagation.

ADELIA MCCREA

RESEARCH DEPARTMENT,  
PARKE, DAVIS AND COMPANY

#### BOOKS RECEIVED

- FOWLER, SIR JAMES K. *The Sthenics*. Pp. 81. Macmillan. \$1.40.
- KINSEY, ALFRED C. *The Gall Wasp Genus Cynips*. Pp. 577. 429 figures. Indiana University Press.
- National Research Council, Committee Report on the Construction and Equipment of Chemical Laboratories*. Pp. xiii + 340. 124 figures. Chemical Foundation.
- PEARSON, H. H. W. *Gnetales*. Pp. vi + 194. 89 figures. Macmillan. \$6.00.
- RILEY, W. A., and R. O. CHRISTENSON. *Guide to the Study of Animal Parasites*. Pp. xv + 131. 32 figures. McGraw-Hill. \$1.50.
- ZELENY, ANTHONY. *Elements of Electricity*. Pp. xxiii + 438. 266 figures. McGraw-Hill. \$3.00.

<sup>1</sup> A. McCrea, SCIENCE, 67, No. 1732, 1928; 69, No. 1798, 1929.



## PURE PROTEINS OF MILK VS. CRUDE CASEIN

**T**HE PURE PROTEINS of milk are so difficult and expensive to prepare, that only too often the crude products are inadvisedly used in researches of importance. One "crude essential element" of the diet may invalidate all other good features of a costly investigation, thereby disqualifying the deduction of the author.

In recent journals such papers have appeared.

"Crude Casein" of the usual industrial type represents simply an acid or rennet curd and is purely "Glorified Pot Cheese," containing unknown products of milk—in unknown amounts.

### FOR RESEARCHES OF PRECISION

*"Casein - Harris" and  
"Lactalbumin - Harris"  
have been prepared*

Through years of study in the Protein Laboratory of Osborne & Mendel (Yale Univ. & Conn. Expt. Sta.), we have learned how to prepare these two proteins in state of highest molecular purity.

Each protein contains highest nitrogen content, lowest mineral ash, freedom from each other and other substances of cow's milk.

They are made directly from fresh skim milk, which is the only possible way to prepare them in high purity.

Assayed upon growing white rats.

Free from all vitamins.

#### AVERAGE ANALYSIS OF CASEIN—HARRIS

Moisture .....	10.73	Casein (N x 6.38) ..	87.09
Ash .....	.59	Ether-soluble .....	.20
Calcium .....	trace	Nitrogen;	
Nitrogen .....	13.66	water, fat, ash-free	15.44

(cf.—Osborne & Harris, Jr. Am. Chem. Soc., 25-IV, 346)

Also

Purest powdered Medicinal Yeast (standardized for Vitamines "B" (F & G) and Concentrates from this yeast.

Also

Vitamine factors, pure food mixes, deficient foods, salt mix and methods for vitamine assays upon the Norwegian white rat.

**THE  
HARRIS LABORATORIES  
TUCKAHOE NEW YORK**

## School of Medicine Western Reserve University

Cleveland, Ohio

**NEW LABORATORIES AND  
HOSPITALS**

**RESTRICTED CLASSES**

**THOROUGH INSTRUCTION**

**LARGE CLINICAL FACILITIES**

**HIGH STANDARD OF SCHOLAR-  
SHIP**

Admission confined to students having academic degrees and to Seniors in Absentia.

*For information address:*

**THE REGISTRAR**

2109 Adelbert Rd.

**CLEVELAND**

## JOHNS HOPKINS UNIVERSITY SCHOOL OF MEDICINE

The School of Medicine is an Integral Part of the University and is in the Closest Affiliation with the Johns Hopkins Hospital.

### ADMISSION

Candidates for admission must be graduates of approved colleges or scientific schools with two years' instruction, including laboratory work, in chemistry, and one year each in physics and biology, together with evidence of a reading knowledge of French and German.

Each class is limited to a maximum of 75 students, men and women being admitted on the same terms. Applications may be sent any time during the academic year but not later than June 15th.

If vacancies occur, students from other institutions desiring advanced standing may be admitted to the second or third year provided they fulfill the requirements and present exceptional qualifications.

### INSTRUCTION

The academic year begins the Tuesday nearest October 1, and closes the second Tuesday in June. The course of instruction occupies four years and especial emphasis is laid upon practical work in the laboratories, in the wards of the Hospital and in the dispensary.

### TUITION

The charge for tuition for 1930-31 will be \$600 per annum, payable in two installments. There are no extra fees except for certain expensive supplies, and laboratory breakage.

Inquiries should be addressed to the  
**Executive Secretary of the School of Medicine, Johns  
Hopkins University, Washington and  
Monument Sts., Baltimore, Md.**

Graduates in Medicine who satisfy the requirements of the heads of the departments in which they desire to work are accepted as students for a period not less than three quarters. Tuition charge is \$50 a quarter.

## School of Medicine and Dentistry THE UNIVERSITY OF ROCHESTER

Medical School, Strong Memorial Hospital, School of Nursing and Out-Patient Department of the University of Rochester and the Municipal Hospital of the City of Rochester, all under one roof. Medical, Surgical, Obstetric, Pediatric, Dental, Contagious and Neurological patients admitted. Unusual opportunities for school and hospital co-operation in medical and graduate dental teaching.

### Admission

Medical candidates must have completed three years of college work with special requirements in chemistry, physics and biology. The entering class will not exceed 50, men and women being admitted on equal terms.

### Tuition

Charge for tuition will be \$400 per annum, payable in equal installments at the beginning of each semester.

*For information address*

**THE DEAN**

School of Medicine and Dentistry  
Crittenden Boulevard, Rochester, N. Y.

## BOSTON UNIVERSITY SCHOOL OF MEDICINE

ORGANIZED IN 1873

### ANNOUNCEMENT

may be obtained by application to

**THE REGISTRAR**

80 East Concord Street,

Boston

Massachusetts

**Allegany School of Natural History**

In Allegany State Park, Quaker Bridge, N. Y.

Fourth Season—July 5 to August 23, 1930

*Registration limited to fifty*

Field studies in Botany, Zoology, Geology, Birds

Natural conditions favorable in richness and variety, suitable laboratories and equipment, comfortable living in a stimulating climate, guidance from experienced teachers and investigators.

Sponsored by the Buffalo Society of Natural Sciences, the New York State Museum, and the University of Buffalo (with college credit).

For circulars or registration, address until June 15:

DR. ROBERT E. COKER, Director  
Chapel Hill, N. C.

New Impression Now Ready

**The Foundations of Science**

By H. POINCARÉ

Pp. xi + 553.

Containing the authorized English translation by George Bruce Halsted of "Science and Hypothesis," "The Value of Science" and "Science and Method," with a special preface by Poincaré, and an introduction by Josiah Royce. *Price, postpaid, \$5.00.*

THE SCIENCE PRESS

Grand Central Terminal

New York, N. Y.

**The New York  
Homeopathic Medical  
College and  
Flower Hospital**

Offers opportunity for highest grade medical training to one hundred men and women selected competitively.

In addition to thorough training in the Preclinical Sciences and the usual courses in Medicine and Surgery, special emphasis is given to Homeotherapy.

Seventy first annual announcement mailed to those interested.

450 East 64th Street  
New York City**The Journal of General Physiology**

EDITED BY

W. J. CROZIER

JOHN H. NORTROP

W. J. V. OSTERHOUT

Contents of Volume 13, No. 4, March 20, 1930:

IRWIN, MARIAN. Spectrophotometric studies of penetration of dyes. VII. Dyes absorbed by a non-medullated nerve.

RAHN, OTTO. The non-logarithmic order of death of some bacteria.

KRUEGER, A. P., and RITTER, R. C. The preparation of a graded series of ultrafilters and measurement of their pore sizes.

CASTLE, E. S. Phototropism and the light-sensitive system of *Phycomyces*.

BETHE, ALBRECHT. The permeability of the surface of marine animals.

DAMON, E. B., and OSTERHOUT, W. J. V. The concentration effect with *Valonia*: Potential differences with concentrated and diluted sea water.OSTERHOUT, W. J. V., and HILL, S. E. Negative variations in *Nitella* produced by chloroform and by potassium chloride.

ANSON, M. L., and MIRSKY, A. E. Protein coagulation and its reversal. The preparation of insoluble globin, soluble globin and heme.

MIRSKY, A. E., and ANSON, M. L. Protein coagulation and its reversal. Improved methods for the reversal of the coagulation of hemoglobin.

BARRON, E. S. GUZMAN, and HOFFMAN, L. A. The catalytic effect of dyes on the oxygen consumption of living cells.

BLINKS, L. R. The direct current resistance of *Nitella*.*THE JOURNAL OF GENERAL PHYSIOLOGY, founded in 1918 by Dr. Jacques Loeb, contains practically all the investigations of Dr. Loeb after that time.**There are, in addition, contributions by many leading physiologists, both in this country and abroad. It has now been decided to offer a complete set of the back volumes at half price to new subscribers.**The first twelve volumes complete may therefore be obtained for \$30.00 by new subscribers until June 30, 1930.*

SUBSCRIPTION PRICE PER YEAR (ONE VOLUME), \$5.00.

PUBLISHED BIMONTHLY BY

**The Rockefeller Institute for Medical Research**  
YORK AVENUE AND 66TH STREET  
NEW YORK, N. Y.