

posed with the formation of ammonium. We have observed no marine bacteria which require urea for their growth.

CLAUDE E. ZOBELL

CATHARINE B. FELTHAM

SCRIPPS INSTITUTION OF OCEANOGRAPHY  
UNIVERSITY OF CALIFORNIA  
LA JOLLA, CALIF.

### A GROWTH-INHIBITING SUBSTANCE IN LETTUCE SEEDS<sup>1</sup>

LETTUCE seeds which fail to germinate on moist blotters at 25° C. in the light may be germinated at this temperature by placing the seeds on moist absorbent cotton or in water.<sup>2</sup> The increased germination obtained by this method suggests that the promotion of germination may be due to the exit of an inhibiting substance which diffuses from the seed into the aqueous medium. If an inhibiting substance is formed during the process of germination then the repeated placing of seeds in contact with the same substratum should cause a gradual reduction in the percentage of germination. The germination tests that have been made on cotton and in water to test this hypothesis show that lettuce seeds do form a substance of unknown nature which diffuses from the seed, and if present in sufficient quantities prevents germination.

The inhibiting material is formed most readily by freshly harvested seeds of the white-seeded varieties of lettuce which tend to go into dormancy at 25° C. Big Boston lettuce seeds which germinated 3 per cent. on moist blotters at 25° C. in the light germinated 80 per cent. when placed on moist cotton. The percentage of germination upon this same cotton medium was reduced to 5 per cent. after five lots of 100 seeds each had been in contact with the medium over a period of 10 days. In like manner germination in a shallow layer of water was completely inhibited after 600 seeds had been in contact with the medium. When the water from a similar inhibiting medium was used to moisten a freshly prepared cotton substratum the germination of lettuce seeds upon the cotton was reduced from 80 per cent. to 10 per cent. A saturated medium which inhibited the germination of Big Boston seeds at 25° C. failed to prevent the germination of Black Seeded Simpson seeds of the same age, which indicates that the physiological condition of the seed is a factor in determining the response made by seeds to the inhibiting substance.

The increased germination of lettuce seeds in the light indicates that light may promote the diffusion

<sup>1</sup> Approved by the director of the New York State Agricultural Experiment Station for publication as Journal Paper No. 53, October 24, 1934.

<sup>2</sup> A. L. Shuck, *New York State Agr. Exp. Sta. Tech. Bul.* No. 222, 1934.

of the substance from the seeds, and although light may accelerate the process, tests have shown that an inhibiting substance passes from the seeds in total darkness. A cotton medium upon which 600 new crop lettuce seeds had been in contact for a period of 10 days in the dark, and then used as a substratum for germination in the light completely inhibited the germination of one-year-old Big Boston seeds. When this cotton medium was washed in water and then used as a substratum a similar lot of seeds germinated 98 per cent.

The age or more specifically the physiological condition of the lettuce seed is a factor influencing the formation of the inhibiting substance, and is also a factor in determining the response made by seeds when placed in contact with a saturated substratum. The inhibiting substance is formed most abundantly by seeds immediately after harvest and in smaller amounts or not at all in old seeds, and appears to be in some way associated with the dormant condition which develops in the seeds when placed at unfavorable temperatures for germination. The marked increase in germination obtained in the light indicates that light may facilitate the passage of the inhibiting material from the seed. The response to light is complicated by the fact that the dormant condition in light-sensitive lettuce seeds can be broken by placing the seeds in an atmosphere that is saturated with water vapor and giving them an exposure to light. The latter response may take place in swollen seeds within a few seconds and without the presence of water in the form of a film surrounding the seed which precludes the possibility of any substance diffusing from the seed. The fact that the material may pass from the seed in total darkness indicates that the function of light is to prevent or break the stable condition of unknown nature which characterizes seeds in secondary dormancy.

A. L. SHUCK

NEW YORK STATE AGRICULTURAL  
EXPERIMENT STATION

### BOOKS RECEIVED

- Carnegie Foundation for the Advancement of Teaching. *Twenty-ninth Annual Report*. Pp. 167. The Foundation, New York.
- Committee on Community Dental Service of the New York Tuberculosis and Health Association. *Health Dentistry for the Community*. Pp. xiii + 85. University of Chicago Press. \$1.00.
- FRITSCH, F. E. *The Structure and Reproduction of the Algae*. Vol. I. Pp. xvii + 791. 245 figures. Macmillan. \$8.00.
- GETTLER, LOTHAR. *Grundriss der Cytologie*. Pp. viii + 295. 209 figures. Von Gebrüder Borntraeger, Berlin. 21.RM.
- Report for the Year 1933-34 of the Department of Scientific and Industrial Research*. Pp. 192. His Majesty's Stationery Office, London. 3s.

# ISO-PROPYL ALCOHOL

## *in Analytical Procedure*

THE use of iso-propyl alcohol in place of ethanol for analytical work is rapidly gaining favor. The higher alcohol has similar properties and possesses the further advantage of being free from restrictions governing its use.


Ferner and Mellon, in *IND. ENG. CHEM.*, (ANAL. ED.) 6, 345 (1934), report new data on solubilities which

indicate wider application of this alcohol in many fields.

Eastman Iso-Propyl Alcohol is supplied in two grades—98-99%, suitable for analytical work, and 90-91%, satisfactory for washing and drying. A resumé of this recent report will be forwarded upon request. Eastman Kodak Company, *Chemical Sales Division*, Rochester, N. Y.



## EASTMAN ORGANIC CHEMICALS




Entrance to the Mead Johnson Research Laboratory, Evansville, Ind. The building is designed to provide for present undertakings and allow for change and expansion required by new projects. Equipment is the best obtainable, modern and adequate. Resources are ample for initiating and conducting investigations of worth.

### *Realizing the importance of original investigations*

in the field of nutrition, Mead Johnson & Company are constantly enlarging their research department. ¶ The results have far surpassed expectations, not only because of the advantages accruing to the Company, *but in contributions* to pure science that have been made. Numerous materials previously unobtainable have been made *available* to scientists at a price only *slightly above* actual cost. ¶ It is a matter of much satisfaction that, without profit,

**Mead Johnson and Company**  
is able to offer assistance to fellow investigators



# ZEISS

## PHOTO-ELECTRIC RECORDING PHOTOMETER

Unexcelled sensitivity

Operates in lighted room

Recording on paper or on plates

Speed of recording 4 cm a minute

Electrometer with bright field and with dark field illumination

Speeds of object and recording surface may be varied continuously  
in ratios from 1:1 to 1:500

Photo-Electric Recording Photometer with motor for 110 volts D. C.  
\$4,936.80 f.o.b. N. Y.

Attachment for darkfield illumination \$174.24

Literature on request

CARL ZEISS, INC.

485 Fifth Avenue  
NEW YORK

728 So. Hill Street  
LOS ANGELES

