

inoculations were successful in 50 per cent. of the attempts. The controls were negative in over 90 per cent. of the patients. These results seem to indicate that *Pityrosporum ovalis* may produce seborrheic dermatitis under favorable conditions.

MORRIS MOORE  
ROY L. KILE

THE BARNARD FREE SKIN AND  
CANCER HOSPITAL  
ST. LOUIS, MISSOURI

#### A DWARF MUTATION IN THE RAT

IN rodents genetic dwarfs have been reported in the guinea pig,<sup>1,2</sup> in the mouse by Snell<sup>3</sup> and recently in the rabbit by Green, Hu and Brown.<sup>4</sup> The dwarfism in the mouse was shown by Smith and MacDowell<sup>5</sup> to be caused by a hereditary deficiency of the anterior pituitary. Preliminary studies in the rabbit suggest that the defect may be due to an endocrine abnormality, while in the guinea pig the nature of the dwarfism was not determined.

The dwarf mutation in the rat described herein was first observed in our colony in the summer of 1933. It appeared as one individual in a litter of nine young in a strain of rats that had been closely inbred in our laboratory for several years. Shortly afterward another dwarf was observed in a litter from a closely related pair, and subsequently in the litters of other closely related rats. At present 22 dwarfs have been produced in 12 litters, the ratio of normal to dwarf in these litters being 80 to 22. These results suggest that the dwarf is the result of a simple autosomal recessive mutation, the appearance of both dwarf males and females showing that the gene is not sex-linked.

At birth normal and dwarf rats can not be distinguished, and it is impossible to separate the dwarf rats with certainty until about the twelfth day after birth. At this time a distinct difference in rate of growth becomes manifest, and the hair of the dwarfs appears much thinner and of finer texture than the hair on their normal litter mates. This difference in the hair is a characteristic feature of the dwarfs throughout life. The mature weight of the dwarfs is approximately 50 per cent. that of the normal males and 70 per cent. of the normal females of this strain. A remarkable feature is the failure of the males and females to become differentiated in size, whereas the male normally becomes distinctly larger than the female.

Thus far all dwarfs have proven to be sterile. No sexual activity has been observed in dwarf males and in only one instance has copulation been noted among the dwarf females. The size of the testes remains infantile, becoming less than one-half normal size. Spermatogenesis occurs, but at a greatly reduced rate, and the few sperm observed were abnormal in appearance. As yet studies have not been made of the ovary.

In all cases so far observed a distinct opacity of the lens of the eyes has been associated with the dwarfism, although some variation in the degree of this opacity exists. Skeletons of the dwarfs have not been prepared, but the general proportion of the body parts appears the same as in normal rats. The effect seems to be a general reduction in size of all parts, and the retardation in rate of growth becomes manifest at a very early date. Using the method of analysis of variance on the weights of litters in which dwarfs appeared, it has been determined that the retardation in growth had begun on the fifth day. As pointed out above, however, individual dwarfs can not be separated with certainty until about the twelfth day after birth. The dwarf rats are weaker than normals, are more susceptible to infections, and are shorter lived.

Individuals heterozygous for the dwarf gene are normal in appearance, in vitality and they attain the same size as homozygous normal rats. Recently, however, some individuals have been observed among the progeny of dwarf-producing pairs that were slightly slow in developing their hair and whose early growth seemed somewhat retarded. Tests are now under way to determine whether such individuals are heterozygous for the dwarf gene.

The decreased size of the dwarf rats, their sterility and their general appearance suggests that the defect may be of endocrine origin. It is, however, apparently not due to a pituitary deficiency, since in preliminary tests with the implantation of pituitary bodies from normal rats, following the technique of Smith and MacDowell, growth was not produced in treated dwarfs. Further experiments to determine the physiologic cause of this dwarfism are at present under way and the colony is being expanded in order to insure a constant supply of the dwarfs.

W. V. LAMBERT  
A. SCIUCHETTI

IOWA STATE COLLEGE

<sup>1</sup> I. J. B. Sollas, Repts. Evol. Com. Roy. Soc., 5: 51-79, 1909.

<sup>2</sup> I. J. B. Sollas, *Jour. Gen.*, 3: 201-204, 1914.

<sup>3</sup> G. D. Snell, *Proc. Nat. Acad. Sci.*, 15: 733-734, 1929.

<sup>4</sup> H. S. N. Green, C. K. Hu and W. H. Brown, *SCIENCE*, n. s., 79: 487-488, 1934.

<sup>5</sup> P. E. Smith and E. C. MacDowell, *Anat. Rec.*, 46: 249-257, 1930.

#### BOOKS RECEIVED

CAULLERY, MAURICE. *French Science*. Pp. xi + 229. Illustrated. Author.  
EDDINGTON, SIR ARTHUR. *New Pathways in Science*. Pp. x + 333. 4 plates. Macmillan. \$3.00.

# THE JOURNAL OF NUTRITION

Published monthly by The Wistar Institute  
John R. Murlin, Managing Editor, University of Rochester

**Vol. 9.**

*Contents for April 1935*

**No. 4.**

- D. B. SMUTS. The relation between the basal metabolism and the endogenous nitrogen metabolism, with particular reference to the estimation of the maintenance requirement of protein.
- AGNES FAY MORGAN, MARION J. HUNT and MILDRED SQUIER. The vitamin B and G content of prunes. One chart, one figure.
- HELEN M. COPLAN and MYRA M. SAMPSON. The effects of a deficiency of iodine and vitamin A on the thyroid gland of the albino rat. Two plates.
- INEZ D. MASON and LEROY S. PALMER. Utilization of gelatin, casein and zein by adult rats.
- MARGARET DANN and GEORGE R. COWGILL. The vitamin C requirement of the guinea pig. One chart.
- E. O. WHITTIER, C. A. CARY and N. R. ELLIS. The effects of lactose on growth and longevity. Five charts.
- SIDNEY FEYDER and H. B. PIERCE. Rates of absorption and glycogenesis from various sugars. Three figures.
- SIDNEY FEYDER. Fat formation from sucrose and glucose.

Price, \$5.00 per volume, Domestic; \$5.50 per volume, Foreign

Two volumes issued annually

Back volumes from 1 to 8, inclusive, may be had for \$40.00

*Address subscriptions to*

**The Wistar Institute of Anatomy and Biology**  
Philadelphia, Pa., U. S. A.

**GRADUATE ASSISTANTSHIPS IN PHYSIOLOGY**—Open to students working toward an advanced academic degree. \$450 with remission of tuition.

*Address:* Chairman, Department of Physiology, Ohio State University, Columbus, Ohio.

**GENERATION OF THE UNIVERSE**  
And "Design For Living"

**The World As Necessity**

Price \$1.50 Postpaid

By PERCY A. CAMPBELL

16208 Nelaview Rd., E. Cleveland, Ohio.

**THE SCIENCE PRESS**  
**PRINTING COMPANY**

PRINTERS OF

SCIENTIFIC AND EDUCATIONAL  
JOURNALS, MONOGRAPHS  
AND BOOKS

*Correspondence Invited*

LANCASTER, PENNSYLVANIA

## BOSTON UNIVERSITY SCHOOL OF MEDICINE

ORGANIZED IN 1873

ANNOUNCEMENT


may be obtained by application to

THE REGISTRAR

80 East Concord Street

Boston

Massachusetts



# ZEISS

## Physicians' and Laboratory Microscope E S C-106

Fixed stage, 12 cm. square  
Illuminating apparatus with  
rack and pinion  
Condenser 1.2 with iris  
Triple revolving nosepiece  
Fine adjustment with gradu-  
ated drum

Achromatic objectives:

8 n.a. 0.20

40 n.a. 0.65

90 n.a. 1.25 oil imm.

Huygens oculars 7x and 10x

Price \$167.50 f. o. b. N. Y.

A good dark-field outfit is obtained by adding: Cardioid condenser \$27.00  
extra for oil im., with iris \$4.00. Compensating ocular 15x: \$12.00

A satchel type of carrying case can be supplied instead  
of the standard cabinet, at an additional cost of \$4.50.

## CARL ZEISS, INC.

485 Fifth Avenue  
NEW YORK

728 So. Hill Street  
LOS ANGELES

