

$$\frac{f_0}{f} = \frac{\rho^{2/3}}{\sqrt{1-\rho^2}} \log_e \frac{1 + \sqrt{1-\rho^2}}{\rho}$$

where $\frac{f_0}{f}$ is the reciprocal of the dissymmetry constant and ρ corresponds to $\frac{a}{b}$ of Kuhn's equation. Substituting the value $1/35$ for ρ , we find a value of 0.396 for $\frac{f_0}{f}$ or 2.52 for $\frac{f}{f_0}$. The molecular weight can be calculated by use of the equation⁹

$$\frac{f}{f_0} = \frac{M \sqrt{(1-Vd)}}{S_{20}} \frac{1}{6\pi\eta N \left(\frac{3}{4}\frac{MV}{\pi N}\right)^{1/3}}$$

M is the molecular weight of the protein, V is the specific volume of the protein in solution, d is the density, S_{20} is the sedimentation constant at 20° C. taken to be 174×10^{-13} ,¹⁰ η is the viscosity coefficient, and N is Avogadro's constant. A value of about 42.5×10^6 , more than 2 times that suggested originally, is found for the molecular weight of the protein by this method. This would correspond to a particle 12.3 μ in diameter and 430 μ in length. This value of the molecular weight is reliable only to the extent to which the equation of Kuhn is applicable to the system under investigation and to which the assumption of no hydration is valid. This treatment of the subject emphasizes the necessity of knowledge of the shape and state of hydration of the tobacco mosaic virus protein in order to enable one to interpret accurately the data from the ultracentrifuge.

MAX A. LAUFFER

THE ROCKEFELLER INSTITUTE FOR
MEDICAL RESEARCH,
PRINCETON, N. J.

THE PRESENCE OF RARE EARTHS IN HICKORY LEAVES

It has been established in the Bureau of Chemistry and Soils that the leaves of the hickory and sweetleaf may contain as much as 1.5 and 6.5 per cent. of crude Al_2O_3 , respectively, in the air-dry leaf. The quantities present in the hickory vary from a few hundredths of 1 per cent. in neutral soils to the above quantities found in acid soils.

The abnormal behavior of the crude alumina precipitate obtained from the hickory leaves from a tree growing in a pegmatite vein of the Moorefield mine, Amelia, Va., led to the separation of a concentrate of the rare earth group of oxides amounting to 0.2 per cent. of the dry weight of the leaves. This figure is probably low, as there may have been some losses in the separations. The colors of the solutions, oxides and oxalates and behavior of the hydroxides indicated

a mixture of cerium, lanthanum, praseodymium and neodymium.

A spectroscopic examination was made at the National Bureau of Standards for the individual rare earths, which are extremely difficult to separate when present in small quantities. The presence and relative abundance of the rare earth elements are shown in the following table:

TABLE 1
SPECTROSCOPIC EXAMINATION OF RARE EARTHS FROM
HICKORY LEAVES*

Element	Oxides from oxalate, several precipitations	Oxides from fluorides, through hydroxides
Cerium	Very strong	Strong
Lanthanum	Strong	Weak
Praseodymium	Strong	Strong
Neodymium	Strong	Strong
Yttrium	Strong	Strong
Samarium	Moderate	Moderate
Europium	Moderate	Moderate
Gadolinium	Very weak	Weak
Dysprosium	Very weak	Weak
Erbium	Trace	Very weak
Ytterbium	Trace	Very weak

* The strengths of the lines of the respective elements were compared using the following scale: Very strong, Strong, Moderate, Weak, Very weak, and Trace.

Scandium, terbium, holmium, thulium and lutecium were not found. Due to lack of information concerning the spectrum of illinium (61), a test was not made for this element.

The presence of "moderate" lines of europium, one of the rarest of the rare earths, is interesting and may point to a concentration of this element by the hickory leaves.

The rare earths are widely distributed and occur in the earth's crust in quantities smaller, but comparable to the quantities of phosphorus and manganese. They have been found in many soils in concentrations up to .05 per cent. The rare earths, lanthanum in particular, are comparatively strong bases. The fact that the rare earths resemble calcium, in forming insoluble oxalates and fluorides, is an indication that the rare earths may substitute for calcium in the growing plant, where that element is deficient. In some of their properties, however, the rare earths more nearly resemble aluminium, and in this case are absorbed by an aluminum-loving plant.

W. O. ROBINSON

RICHARD WHETSTONE

BUREAU OF CHEMISTRY AND SOILS

BOURDON F. SCRIBNER

BUREAU OF STANDARDS

BOOKS RECEIVED

LEWIS, BERNARD and GUENTHER VON ELBE. *Combustion, Flames and Explosions of Gases*. Pp. xiii + 415. 79 figures. Cambridge University Press; Macmillan. \$5.50.

SHULL, A. FRANKLIN. *Heredity*. Third edition. Pp. xvii + 442. 168 figures. McGraw-Hill. \$3.50.

⁹ T. Svedberg, *Chem. Rev.*, 14: 1, 1934.

¹⁰ R. W. G. Wyckoff, *loc. cit.*