



Supplementary Materials for

A Global Pattern of Thermal Adaptation in Marine Phytoplankton

Mridul K. Thomas,* Colin T. Kremer, Christopher A. Klausmeier, Elena Litchman

*To whom correspondence should be addressed. E-mail: thomasmr@msu.edu

Published 25 October 2012 on *Science Express*

DOI: 10.1126/science.1224836

This PDF file includes:

Materials and Methods
Figs. S1 to S11
Tables S1 to S4
Captions for Tables S5 and S6
References (35–135)

Other Supplementary Materials for this manuscript include the following:

(available at www.sciencemag.org/cgi/content/full/science.1224836/DC1)

Fig. S12
Tables S5 and S6

1 Materials and Methods

We provide additional details on methods, statistical analyses, models, and data sources in this supplement. A graphical overview of the various data sets and analyses composing this project can be seen in Figure S11. §1.1 describes the selection criteria used to identify the growth rate data used in this study. §1.2 details the statistical procedures used to fit and describe the thermal tolerance curves. §1.3 outlines the bootstrapping approach used to appropriately account for uncertainty in estimates of thermal traits. §1.4 covers the statistical analysis used to characterize temperature regimes. §1.5 describes statistical methods used to demonstrate trait-environment relationships while accounting for trait uncertainty. §1.6 explains our analysis of trait variation as a function of taxonomy. §1.7 describes the structure and analysis of our eco-evolutionary model. §1.8 provides extra details on the data and methods employed in our species distribution models. §1.9 lists the various pieces of computational software we employed.

1.1 Curve selection criteria

We assembled a data set containing growth rate measurements of marine and estuarine phytoplankton at different temperatures that have been published over the past century. All measurements were digitized using *g3data* (35). Strains were treated as independent, due to the existence of considerable intraspecific variation and uncertainty in species boundaries for some taxa.

Several criteria were used to determine the inclusion of species/strain data in our analyses.

- 1) To facilitate comparisons across studies we only included data for growth rates measured in units that could be converted to specific growth rate.
- 2) Because we were primarily concerned with estimating the temperature at which strains/species achieve their maximum growth rates, we rejected curves where the largest measured growth rate occurred at the lowest or highest temperature considered.
- 3) Curves with fewer than four measured growth rates were excluded, as were curves showing strong bimodality, which we attributed to imprecise experimental measurements.
- 4) Where curves were measured under different experimental conditions (salinity, nutrient limitation, light levels, day length), we preferentially selected curves meeting the following conditions:
 - a. Salinity between 30 and 40 parts per thousand.
 - b. Light levels greater than or equal to 100 microeinsteins.m⁻².s⁻¹.
 - c. Not experimentally limited by nutrients
 - d. Day lengths of greater than or equal to 10 hours.

When no curves for a particular isolate satisfied these experimental constraints, we settled for using data from the curve(s) that were closest to the desired light and salinity levels.

- 5) We considered only marine and estuarine strains not isolated from inland waters.

After applying these criteria, we had data for a total of 252 separate curves, divided among 194 isolates/strains belonging to approximately 130 species, from 111 unique isolation locations ranging in latitude from 76°N to 75°S (Fig. S1, Table S5).

1.2 Statistical analysis of thermal tolerance curves

Temperature dependent specific growth rates can be described by the following equation¹⁶:

$$f(T) = ae^{bT} \left[1 - \left(\frac{T-z}{w/2} \right)^2 \right] \quad (\text{S.1})$$

Here specific growth rate f is an explicit function of temperature, T . The shape of the thermal tolerance curve is controlled by two important species traits, z and w . The range of temperatures over which growth rate is positive, or the thermal niche width, is given by w . Species trait z determines the location of the maximum of the quadratic portion of this function. In the case where parameter $b = 0$, this value is identical to the temperature at which a species achieves its maximum growth rate. However, when b is non-zero, the maximum value of S.1 falls above (or potentially below, $b < 0$) the value of z , and can be found through numerical optimization.

Norberg (14) fixed parameters a and b according to the values of the Eppley curve (12), an exponential relationship thought to provide the community-level upper bound (95% quantile) on phytoplankton growth as a function of temperature. However, in fitting growth curves to data for individual strains, we recognized that species may not strictly follow this community level constraint, potentially due to the effects of other constraints such as light or nutrient limitation. For this reason, we allowed a and b to be free parameters, fit simultaneously with z and w .

To describe the growth data for each isolate, we used a maximum likelihood approach, such that the mean growth rate at a given temperature followed equation (S.1),

$$\mu = f(T) + \mathcal{N}(0, \sigma^2) \quad (\text{S.2})$$

Here observational error was described by a normal distribution with a mean of zero and variance of σ^2 .

1.3 Determining physiological parameter uncertainty

While confidence intervals for the point estimates of the parameters of (S.1) were easy to obtain, it was not straightforward to determine uncertainty for implicit properties such as the temperature at which growth rate is maximized (or, the ‘optimum temperature’). Yet, this was the property that we were mainly interested in, leading us to adopt a parametric bootstrapping approach.

We used a Monte Carlo approach such that for each thermal tolerance curve having n data points, we simulated n new data points, drawn from a normal distribution such that:

- 1) The mean of the distribution corresponds to the value of (S.1) at each of the original experimental temperatures, given the coefficients previously estimated for the original curve.
- 2) The standard deviation of the distribution, σ , was obtained by adjusting the original maximum likelihood estimate, $\hat{\sigma}$, to account for uncertainty in its estimation (36):

$$\hat{\sigma}^2 = \sigma^2 \sqrt{(n-1)/X} \quad (\text{S.3})$$

where n is the number of points, and X is a random number drawn from the χ^2 distribution having $(n - 1)$ degrees of freedom.

Equation (S.1) was then fit to the simulated data using maximum likelihood estimation, and the new parameter values, as well as the numerically estimated optimum temperature, were retained. Repeating this process a total of 10,000 times (for each isolate), yielded bootstrapped distributions of all parameter estimates. From these distributions we calculated the 95% confidence intervals as the range between the 2.5th and 97.5th quantiles. These estimates of uncertainty were vital for subsequent analyses, as they allowed us to appropriately account for the inherent differences in uncertainty between different isolates expected to arise any time that data are synthesized across many individual studies.

1.4 Environmental data analysis

To provide the environmental data necessary for investigating trait-environment relationships, we turned to historical sea surface temperature (SST) estimates available through NOAA as ¼ degree, AVHRR/AMSR+AVHRR daily optimum interpolation SST (18). These data covered the period between 1981 and 2010. Because of the fine spatial resolution of these data, we were able to closely match the location of each isolate to the nearest location having SST data, minimizing error due to spatial variation. When more than one ¼ degree location was equidistant from an isolation location, a specific grid location was selected randomly from the various options. Our initial set of 111 distinct isolation locations were matched in this manner to a set of 106 distinct SST locations. We then assembled time series of sea surface temperature at each of these locations from Sept. 1st, 1981 to Jan. 18th, 2011. Temperature regimes were described by fitting the following modified sinusoidal function to the entire 30-year time series at each location, again using a maximum likelihood approach:

$$T(t) = \phi + r \left[\sin \left(\frac{\pi}{365} (t + \beta) \right) \right]^{2\alpha} \quad (\text{S.4})$$

In this model, $|r|$ describes the range of temperatures achieved, while ϕ provides the maximum ($r < 0$) or minimum ($r > 0$) temperature. Parameters α and β describe the skewness and temporal shift of the temperature oscillations, respectively. The addition of the α parameter extends a typical sine function to capture asymmetrical seasonality (for example, longer warm periods than cold periods). Note however, that for $\alpha = 1/2$, equation (S.4) reduces to an ordinary sinusoidal model. These regressions generally fit the data well ($n = 106$ time series, mean R^2 of 0.81, standard deviation of 0.17). Note that due to

the inherent assumption of observation error, standard regression models describing temperature time series (such as S.4) will always underestimate the extreme range of temperatures.

We calculated mean temperatures and temperature ranges directly from time series data for each location, as well as the mean and range of the deterministic model fit to the time series. These results are used subsequently in establishing trait-environment relationships, and (in the case of the deterministic model) to describe realistic temperature fluctuations in our eco-evolutionary model.

1.5 Trait-environment regression randomizations

We wanted to calculate regressions relating latitude and mean temperature to phytoplankton optimum temperatures, while accounting for the varying levels of uncertainty attached to each estimate of optimum temperature. Typically meta-analyses handle this issue by performing a weighted regression, using the inverse of the uncertainty associated with each data point. This approach implicitly assumes that the uncertainty around each point estimate is symmetric, which was clearly not the case for our traits (see for example the bootstrapped confidence intervals around point estimates of optimum temperatures, Fig. S9). Consequently, we employed a different approach using resampling techniques, which, to be useful, needed to provide: 1) regression coefficients, 2) uncertainty in regression fit, and 3) a means of comparing the fits of a set of alternative models. Note that we wanted to account for two kinds of uncertainty in this process, both the uncertainty arising from error in estimates of optimum temperature and the uncertainty associated with fitting a regression to data.

Recall that during our earlier parametric bootstrapping calculations we obtained 10,000 estimates of the thermal optimum of each of $n = 194$ isolates, via a process that is independent across isolates. For the i th iteration of this resampling procedure, we did the following:

- 1) Drew the i th value (out of 10,000) from each of the 194 distributions of estimated thermal optima. This captured error associated with uncertainty in our thermal optima estimates.
- 2) Fit each of the regression models under consideration to this randomized data
- 3) Performed standard model comparison of these regressions, saving the Akaike weights associated with each model. Akaike weights capture the relative likelihood of a model, given the data and set of competing models under consideration (37).
- 4) To capture uncertainty associated with each of these models, we conducted residual resampling, associating each predicted value from a given regression with a new residual, and repeating the regression.
- 5) Saved the estimates of the resulting regression coefficients.

This process was repeated a total of 10,000 times across all of the bootstrapped optimum temperature values. The result was a distribution of parameter estimates for each of the regression fits. From these we calculated confidence intervals on the regression

parameters, as well as confidence bands for the relationships as a whole. Finally, we determined the model(s) with the most explanatory power by looking at the average Akaike weight (across the 10,000 replicates) associated with each.

This method was used to examine the relationship between optimum temperature and latitude (having considering linear and quadratic models), resulting in Figure 1. A more extensive set of models was examined when considering the relationship between optimum temperature and the mean and range of environmental temperatures. The set of models considered, as well as the corresponding average Akaike weights, are provided in Table S4, while the model receiving the greatest weight is presented in Figure 2A.

1.6 Taxonomic model comparison

As the genetic sequence data necessary to construct a complete phylogeny for the species in our dataset were not available, we tested for a taxonomic signal (38, 39) in optimum temperature and temperature niche width. Strains were classified according to Algaebase (40) supplemented with a few entries from ITIS (41). In the case of optimum temperature, we constructed a full linear mixed model containing environmental parameters (mean annual temperature) as fixed effects and all taxonomic levels (from domain to strain) as nested random effects. This was compared against a set of models that omitted each individual taxonomic level using Akaike Information Criterion (AIC), a tool used in information theoretic approaches to model selection that measures goodness of fit of a statistical model (Table S1). In the case of niche width, the models contained taxonomic levels, but not environmental covariates, as none explained variation in niche width. A summary of the full models is in Table S2. Analyses were repeated using an alternate method in which taxonomic levels were added in sequentially and compared against the previous model. The results did not change our conclusions, though we were unable to test for the effect of Domain on either trait.

1.7 Eco-evolutionary modeling

1.7.1 Model setup.

We used a version of an eco-evolutionary modeling framework called adaptive dynamics (15, 16) to determine the thermal strategy or strategies that represent evolutionary equilibria for each temperature regime represented in our dataset. All parameters used in the subsequent model are presented in Table S3, along with their units and the numerical values used in our simulations, where appropriate. Our approach consisted of defining the per capita growth rate of phytoplankton strain i as a function of its traits (focusing on z_i) and environment (nutrient level and temperature) as follows:

$$\frac{1}{N_i} \frac{dN_i}{dt} = \mu(z_i, T) \frac{R}{R+k} - m(T) \quad (\text{S.5})$$

In this model, strains exhibit a temperature dependent growth rate $\mu(z_i, T)$, equation (S.7), subject to the availability of resource R and half-saturation constant k . Given that very

little is known about the interactive effects of temperature and resource dynamics, we treated resource dynamics as simply as possible (algebraically):

$$R(t) = R_{in} - \sum_i N_i(t) \quad (\text{S.6})$$

Growth rates were taken to be both trait and temperature dependent:

$$\mu(z_i, T(t)) = 0.81e^{0.0631T(t)} \left[1 - \left(\frac{T(t) - z_i}{w/2} \right)^2 \right] \quad (\text{S.7})$$

where temperature dynamics follow (S.4). This equation is similar to that of Norberg (14) and (S.1), but instead of using Eppley's original coefficients (12) we made use of more recently refined parameters (42). The value of this function (S.7) depends critically on strain traits z_i and w , which are the temperature at which a strain touches the Eppley curve and the strain's thermal niche width, respectively. Three different values of niche width w were investigated, approximately spanning the range of niche widths that we could estimate from literature data with confidence (6.1 to 33.9 °C). Trait z_i can be interpreted as the temperature at which a strain's thermal tolerance curve lies tangent to the Eppley curve. To obtain predictions of thermal optima comparable to those in the rest of the paper, we used numerical methods to determine the temperature at which (S.7) reaches its maximum. In addition to depending on trait z_i , growth rate depends on the temperature, $T(t)$, which varies over time following (S.4) with specific parameters that differ depending on the particular environment under consideration (see section 'Environmental data analysis'). In the model, all strains experience a temperature dependent mortality rate, $m(T)$ that is 5% of the Eppley curve (i.e. 5% of the maximum growth rate), regardless of their traits.

Finally, we allowed strains to evolve or change their trait value z_i dynamically through time according to (16):

$$\frac{dz_i}{dt} = \varepsilon \frac{d}{dz_i} \left[\frac{1}{N_i} \frac{dN_i}{dt} \right] \quad (\text{S.8})$$

In this model, the change in trait over time is driven by the fitness gradient of a strain, or how much the per capita growth rate (fitness) of a strain increases (or decreases) with small changes in its trait value. This rate is scaled by parameter ε , which we take to be small so that traits are approximately constant within a period, effectively forcing the separation of time scales between ecological and evolutionary dynamics (43). Equation (S.8) is analogous to the breeder's equation drawn from quantitative genetics (16).

1.7.2 Predictions of optimal trait values - adaptive dynamics simulations.

Together, (S.5-S.8) provide a system of differential and algebraic equations that can be analyzed numerically and used to predict the evolutionary stable strategies (ESSs), or equilibrium trait value(s), of phytoplankton species in a given environment. These

predictions allowed us to test whether our understanding of the interaction between physiological constraints and environmental variation, and ecology and evolution, is enough to predict trait-environment relationships observed in the real world. Eco-evolutionary models have rarely been used as a source of predictions for comparing with data, with notable exceptions (17, 44, 45).

We performed eco-evolutionary simulations for each of the 106 distinct historical temperature regimes earlier matched to isolates. The temperature forcing (S.4) for these simulations are parameterized using the results from the earlier section ‘Environmental data analysis’. For each location, we started with four species having initial trait values z_i uniformly distributed across that site’s range of temperatures. We then used Wolfram Mathematica version 8 to numerically solve this system of eight differential equations forward in time, until they reached their dynamic attractor. During this process, the trait and population dynamics of each strain were monitored, such that:

- 1) If the trait values of any two strains converged on each other to within 1×10^{-4} , their biomasses were summed, and they were merged into a single pair of equations, and
- 2) If the biomass of an individual strain integrated over one time period fell below 1×10^{-12} , it was considered to have suffered extinction and its pair of equations was removed from the system.

Once this system reached its dynamic attractor, we took the average value of z_i over a single period for each of the remaining strains and determined the temperature at which the corresponding thermal tolerance curve achieved its maximum growth rate. This provided us with estimates of the ESS optimum temperatures and diversity of coexisting strains, matched to that particular environment (Fig. S10).

We then repeated this process across each set of environmental conditions, for three different settings of niche widths (see Fig. S3 and parameters in Table S3). For environments where the ESS consisted of a single strain, we found that its optimum temperature was consistently higher than the mean annual temperature by a factor determined by the strain's niche width. These strains all shared the property that their z (the temperature at which their thermal tolerance curves were tangent to the Eppley curve) exactly matched the mean annual temperature they experienced. In other words, their most adaptive strategy is to have values of z such they grew as fast as theoretically possible at the mean annual temperature. However, because their thermal tolerance curves are bounded by an increasing function, the value of z is not the same as the temperature at which each strain achieves its own fastest growth rate, usually defined as its optimum temperature (see Fig. S2). The difference between the temperatures at which these two points occur is determined by the niche width, explaining the offset we observe.

This particular method of eco-evolutionary modeling is only one of several methods that could be applied to this problem. We selected it because under some situations, such as with fluctuating environments, it can be more computationally tractable than more traditional ‘Adaptive Dynamics’ methods. Additionally, in future

work it can easily be extended to account for trait change on ecological time scales (16, 46) by increasing ε , enabling us to potentially study evolutionary responses to climate change within the same framework. It is worth noting that the maximum number of coexisting strains that can be identified using this approach is constrained by the total number of initial strains (four in our case), so it remains possible that temperature variation could support higher levels of diversity than we determined in this study. However, this seems unlikely as the most diverse result obtained consisted of only three strains.

1.8 Species distribution modeling.

The thermal tolerance curve of a given species allowed us to calculate its growth rate under any given environmental temperature. A time series of environmental temperatures can thus be converted into a time series of estimated growth rates. We can then calculate the average per-capita growth rate of a species over the duration of the time series, in the absence of other limiting factors. Environments in which this growth rate was positive were considered to be within the geographic limits of the species' fundamental niche, with respect to temperature. Repeating this exercise for time series data across the world oceans for each strain in our study, we estimated the spatial extent of the strain's fundamental temperature niche. We then examined how these species distributions differ given historical and predicted future thermal regimes.

1.8.1 Data sets

For this analysis we drew on two additional sources of sea surface temperature data:

- 1) Monthly mean historical sea surface temperatures, $1^\circ \times 1^\circ$ (19), and
- 2) Projections of future temperature regimes obtained from the NOAA GFDL CM2.1 (23, 47). This model was driven with the SRES A2 emissions scenario, characterised by rapid population growth and heterogeneous development (22). This dataset contained global ocean temperature projections from 2001 to 2100, with a spatial resolution of $1^\circ \times 1^\circ$ and a temporal resolution of 1 month. We focused on the final ten-year period from 2091-2100, at the end of which CO_2 concentrations reach 800 ppm.

These data sets have similar resolutions, but use different spatial grids (fixed versus Gaussian, respectively). To generate predictions that are directly comparable between historical and future temperature regimes we used bounded universal kriging to conduct spatial interpolations, basing the requisite theoretical variograms on Gaussian model fits to the semivariograms of each spatial data set. Because the world is round, kriging methods applied to global data sets based on latitude/longitude are prone to boundary effects. We minimized this problem for our kriging by generating two different interpolations, where the longitudinal boundary fell at 0° and 180° , respectively. We then generated a corrected interpolation by combining these results, while excluding edge values within the local neighbourhood size employed in the bounded universal kriging. This interpolation allowed us to generate spatially congruent data sets, based on which we could make valid area, range shift and diversity comparisons.

1.8.2 Range shift estimates.

Using the above methods, we generated spatial predictions of the fundamental temperature niches (i.e. all locations where its growth rate averaged over its thermal environment was positive) or species distributions of all 194 strains in our data set, for both historical and future environments. These calculations were performed without allowing for strain evolution, such that the traits of contemporary strains were assumed to be the same traits with which they will confront future environments. Comparing these results enabled us to quantify shifts in the fundamental range of these species, including changes in both range size and range position (see Figs. S4–S6, and Fig. S12 for a complete set of range maps).

1.8.3 Diversity change estimates.

When examined in the aggregate, species distribution models allowed us to examine broad-scale patterns in potential diversity. We generated potential diversity predictions for each ocean location by calculating the number of strain ranges it fell within, under both historical and predicted future temperature regimes. Historical diversity estimates were kriged (see above) to make them comparable to estimates of future diversity. The historical diversity at each location was then subtracted from future diversity to generate predictions of diversity change over the 100-year time period (see Figs. S7 and S8).

1.9 Software employed.

All statistical analyses were performed using R (48), as well as the following packages: *bbmle* (49) (for maximum likelihood regressions and model comparison), *reshape* (50) (for data manipulation), *ggplot2* (51) (for figure creation), *ncdf4* (52) (for working with environmental data in NetCDF formats), *gstat* (53) (for performing spatial interpolation/kriging), and *lme4* (54) (for mixed model analyses).




Fig. S1.

Mean annual temperatures across the oceans and the isolation locations of the 194 strains in our dataset, indicated by white dots. While most strains are isolated from coastal regions, we capture almost the entire temperature gradient, including the polar and tropical extremes.




Fig. S2.

An example thermal tolerance curve, illustrating the skewness typical of all known ectotherms, including reptiles, amphibians, fish, algae, bacteria, and viruses. Niche width and the optimum temperature (at which the strain reaches its maximum growth rate) are shown. As in our model, this strain is bounded by the Eppley curve.






Fig. S3.

Eco-evolutionary model predictions of the evolutionarily stable optimum temperatures for growth given different niche widths (panels A-C; $w = 10, 20,$ and 30°C , respectively). Also shown in D is the number of environments for which the equilibrium community consisted of 1, 2, or 3 coexisting strains, displayed for each of the various niche width assumptions. As niche width increases the frequency of obtaining multiple coexisting species declines.

A**B****Fig. S4.**

Predicted shifts in the A) equatorial and B) polar boundaries of all 194 strains. A) Points above the 1:1 dashed line have experienced a poleward shift in their lowest latitude at which they can grow. Almost all strains experience no change or poleward shifts. B) Points above the 1:1 dashed line have experienced a poleward shift in their highest latitude at which they can grow. Most strains experience a fairly small change in this latitude.




Fig. S5.

Examples of changes in the fundamental niche illustrating the diversity of ways individual strains may be affected. At each location a strain can persist (grey), be absent from the environment (white), or undergo range expansion (blue) or contraction (red) in the future. See figure S12 for the complete set of range maps.




Fig. S6.

Histogram of per cent change in predicted strain range sizes. A number of strains experience a slight increase in their range, but a large number experience small to moderate decreases.




Fig. S7.

Potential diversity under past (1991-2000) temperature regimes. High diversity in temperate waters is a result of sampling bias, as most strains in our dataset were isolated in temperate waters, specifically off the US east coast, the European west coast and Japan.




Fig. S8.

Potential diversity under future (2091-2100) temperature regimes. Temperature increase drives a large reduction in the potential diversity of the tropical Indian Ocean, Pacific Ocean and western Atlantic Ocean and an increase in the Antarctic Ocean.




Fig. S9.

Estimates of uncertainty in optimum temperature for all strains, obtained through bootstrapping. Estimated optima are shown as points, while the 95% confidence intervals for the point estimates are shown by the error bars, obtained via our randomization analysis. Strains are ranked in ascending order of optimum temperature.




Fig. S10.

Example time series of population dynamics and corresponding species traits. The first row contains results for $w = 20$, while the second shows results for the same environment, but with a narrower niche width ($w = 10$) leading to three coexisting strains.




Fig. S11.

A schematic outlining all analyses and modeling performed.

Table S1.

Taxonomic model comparisons. Taxonomic levels are considered significant if their removal leads to an increase in AIC of >2 (i.e. $\Delta\text{AIC} < -2$) relative to the full model, and are indicated here in boldface.

Trait	Model	AIC	dAIC
Optimum temperature for growth	Full model	1091.407	0
	Full model - Domain	1089.407	2
Full model:	Full model - Kingdom	1090.960	0.447
	Full model - Phylum	1089.407	2
	Full model - Class	1089.407	2
Mean temperature + (Mean temperature) ² + Domain + Kingdom + Phylum + Class + Order + Family + Genus + Species + Strain	Full model - Order	1089.407	2
	Full model - Family	1089.407	2
	Full model - Genus	1092.006	-0.599
	Full model - Species	1098.854	-7.447
	Full model - Strain	1089.447	1.96
	Temperature niche width	Full model	1522.816
Full model:	Full model - Domain	1521.898	0.918
	Full model - Kingdom	1520.816	2
Domain + Kingdom + Phylum + Class + Order + Family + Genus + Species + Strain	Full model - Phylum	1522.256	0.56
	Full model - Class	1520.816	2
	Full model - Order	1520.816	2
	Full model - Family	1520.816	2
	Full model - Genus	1521.336	1.48
	Full model - Species	1546.125	-23.309
	Full model - Strain	1520.816	2

Table S2.

A summary of the full models (as written in Table S1) used to test for a taxonomic signal. Significant random effects indicated in boldface.

Optimum temperature for growth				Temperature niche width			
Random effects				Random effects			
<i>Taxonomic level</i>	<i>Variance explained</i>			<i>Taxonomic level</i>	<i>Variance explained</i>		
Domain	<0.001			Domain	52.044		
Kingdom	0.937			Kingdom	<0.001		
Phylum	<0.001			Phylum	14.775		
Class	<0.001			Class	<0.001		
Order	<0.001			Order	<0.001		
Family	<0.001			Family	<0.001		
Genus	1.867			Genus	7.142		
Species	4.722			Species	55.514		
Strain	0.173			Strain	<0.001		
Residual	8.868			Residual	91.489		
Fixed effects				Fixed effects			
<i>Term</i>	<i>Estimate</i>	<i>Std. error</i>	<i>t-value</i>	<i>Term</i>	<i>Estimate</i>	<i>Std. error</i>	<i>t-value</i>
Intercept	7.216	1.141	6.326	Intercept	14.114	5.875	2.402
Mean temperature	1.358	0.133	10.238				
(Mean temperature) ²	-0.024	0.004	-5.657				

Table S3.

Eco-evolutionary model parameters.

	Symbol	Definition	Value or units
Resource variables	R	Available resource	Dynamic
	k	Half-saturation constant	1
	R_{in}	Total resource level	100
Biomass variables	N	Biomass	Dynamic; measured in resource units
	$m(T)$	Temperature dependent background mortality rate	Specific growth rate; $m = 0.5(ae^{bT})$
	μ	Growth rate	Specific growth rate
	a	Eppley curve coefficient	0.81
	b	Eppley curve exponent	0.0631
	z_i	Competitive optima	°C
	w	Niche width	{10, 20, 30}, °C
Temperature variables	T	Temperature	°C
	τ	Period of temperature fluctuation	365 days
	ϕ	Minimum (maximum) temperature, if r is positive (negative)	°C
	r	Range of temperature fluctuation	°C
	α	Skewness of temperature fluctuation	Not applicable
	β	Temporal shift of temperature fluctuation	Days
Evolutionary parameters	ε	Evolution rate	0.01

Table S4.

Trait-environment models and associated Akaike weights.

Model	Average Akaike weight
Optimum ~ Mean	0.0153
Optimum ~ Range	0.0002
Optimum ~ Mean + Range	0.0313
Optimum ~ Mean + Mean ²	0.6661
Optimum ~ Mean + Mean ² + Range	0.2871

Table S5 (separate file).

Phytoplankton temperature traits and data sources (citations 55-135).

Table S6 (separate file).

Phytoplankton growth rates and isolation locations, gathered from citations 55-135.

References and Notes

1. C. B. Field, M. J. Behrenfeld, J. T. Randerson, P. G. Falkowski, Primary production of the biosphere: integrating terrestrial and oceanic components. *Science* **281**, 237 (1998). [doi:10.1126/science.281.5374.237](https://doi.org/10.1126/science.281.5374.237) [Medline](#)
2. A. C. Redfield, The biological control of chemical factors in the environment. *Am. Sci.* **46**, 205 (1958).
3. P. G. Falkowski, R. T. Barber, V. Smetacek, Biogeochemical controls and feedbacks on ocean primary production. *Science* **281**, 200 (1998). [doi:10.1126/science.281.5374.200](https://doi.org/10.1126/science.281.5374.200) [Medline](#)
4. M. J. Behrenfeld *et al.*, Climate-driven trends in contemporary ocean productivity. *Nature* **444**, 752 (2006). [doi:10.1038/nature05317](https://doi.org/10.1038/nature05317) [Medline](#)
5. D. G. Boyce, M. R. Lewis, B. Worm, Global phytoplankton decline over the past century. *Nature* **466**, 591 (2010). [doi:10.1038/nature09268](https://doi.org/10.1038/nature09268) [Medline](#)
6. M. Edwards, A. J. Richardson, Impact of climate change on marine pelagic phenology and trophic mismatch. *Nature* **430**, 881 (2004). [doi:10.1038/nature02808](https://doi.org/10.1038/nature02808) [Medline](#)
7. X. A. G. Morán, Á. López-Urrutia, A. Calvo-Díaz, W. K. W. Li, Increasing importance of small phytoplankton in a warmer ocean. *Glob. Change Biol.* **16**, 1137 (2010). [doi:10.1111/j.1365-2486.2009.01960.x](https://doi.org/10.1111/j.1365-2486.2009.01960.x)
8. L. Bopp *et al.*, Potential impact of climate change on marine export production. *Global Biogeochem. Cycles* **15**, 81 (2001). [doi:10.1029/1999GB001256](https://doi.org/10.1029/1999GB001256)
9. M. Steinacher *et al.*, Projected 21st century decrease in marine productivity: A multi-model analysis. *Biogeosciences* **7**, 979 (2010). [doi:10.5194/bg-7-979-2010](https://doi.org/10.5194/bg-7-979-2010)
10. Materials and methods are available as supplementary materials on *Science Online*.
11. J. G. Kingsolver, The well-temperated biologist. *Am. Nat.* **174**, 755 (2009). [doi:10.1086/648310](https://doi.org/10.1086/648310) [Medline](#)
12. R. W. Eppley, Temperature and phytoplankton growth in the sea. *Fish Bull.* **70**, 1063 (1972).
13. T. L. Martin, R. B. Huey, Why “suboptimal” is optimal: Jensen’s inequality and ectotherm thermal preferences. *Am. Nat.* **171**, E102 (2008). [doi:10.1086/527502](https://doi.org/10.1086/527502) [Medline](#)
14. J. Norberg, Biodiversity and ecosystem functioning: A complex adaptive systems approach. *Limnol. Oceanogr.* **49**, 1269 (2004). [doi:10.4319/lo.2004.49.4_part_2.1269](https://doi.org/10.4319/lo.2004.49.4_part_2.1269)
15. S. A. H. Geritz, É. Kisdi, G. Meszéna, J. A. J. Metz, Evolutionarily singular strategies and the adaptive growth and branching of the evolutionary tree. *Evol. Ecol.* **12**, 35 (1997). [doi:10.1023/A:1006554906681](https://doi.org/10.1023/A:1006554906681)
16. P. A. Abrams, Modelling the adaptive dynamics of traits involved in inter- and intraspecific interactions: An assessment of three methods. *Ecol. Lett.* **4**, 166 (2001). [doi:10.1046/j.1461-0248.2001.00199.x](https://doi.org/10.1046/j.1461-0248.2001.00199.x)
17. J. C. Stegen, R. Ferriere, B. J. Enquist, Evolving ecological networks and the emergence of biodiversity patterns across temperature gradients. *Proc. Biol. Sci.* **279**, 1051 (2011). [10.1098/rspb.2011.1733](https://doi.org/10.1098/rspb.2011.1733) [Medline](#)

18. R. W. Reynolds *et al.*, Daily high-resolution-blended analyses for sea surface temperature. *J. Clim.* **20**, 5473 (2007). [doi:10.1175/2007JCLI1824.1](https://doi.org/10.1175/2007JCLI1824.1)
19. R. W. Reynolds, N. A. Rayner, T. M. Smith, D. C. Stokes, W. Wang, An improved in situ and satellite SST analysis for climate. *J. Clim.* **15**, 1609 (2002). [doi:10.1175/1520-0442\(2002\)015<1609:AIISAS>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<1609:AIISAS>2.0.CO;2)
20. M. R. Kearney, W. Porter, Mechanistic niche modelling: combining physiological and spatial data to predict species' ranges. *Ecol. Lett.* **12**, 334 (2009). [doi:10.1111/j.1461-0248.2008.01277.x](https://doi.org/10.1111/j.1461-0248.2008.01277.x) [Medline](#)
21. I. P. C. C. Fourth Assessment Report, *IPCC, 2007: Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and (Cambridge, United Kingdom and New York, NY, USA, 2007), p. 996.
22. N. Nakicenovic *et al.*, *Special report on emissions scenarios: A special report of Working Group III of the Intergovernmental Panel on Climate Change* N. Nakićenović, R. Swart, Eds. (Cambridge University Press, Cambridge, UK, 2000; http://www.osti.gov/energycitations/product.biblio.jsp?osti_id=15009867), p. 595.
23. T. Delworth *et al.*, GFDL's CM2 global coupled climate models. Part I: Formulation and simulation characteristics. *J. Clim.* **19**, 643 (2006). [doi:10.1175/JCLI3629.1](https://doi.org/10.1175/JCLI3629.1)
24. D. W. McKenney, J. H. Pedlar, K. Lawrence, K. Campbell, M. F. Hutchinson, Potential impacts of climate change on the distribution of North American trees. *Bioscience* **57**, 939 (2007). [doi:10.1641/B571106](https://doi.org/10.1641/B571106)
25. D. U. Hooper *et al.*, A global synthesis reveals biodiversity loss as a major driver of ecosystem change. *Nature* **486**, 105 (2012). [Medline](#)
26. D. Tilman, D. Wedin, J. Knops, Productivity and sustainability influenced by biodiversity in grassland ecosystems. *Nature* **379**, 718 (1996). [doi:10.1038/379718a0](https://doi.org/10.1038/379718a0)
27. P. B. Reich *et al.*, Impacts of biodiversity loss escalate through time as redundancy fades. *Science* **336**, 589 (2012). [doi:10.1126/science.1217909](https://doi.org/10.1126/science.1217909) [Medline](#)
28. C. A. Deutsch *et al.*, Impacts of climate warming on terrestrial ectotherms across latitude. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 6668 (2008). [doi:10.1073/pnas.0709472105](https://doi.org/10.1073/pnas.0709472105) [Medline](#)
29. K. Härnström, M. Ellegaard, T. J. Andersen, A. Godhe, Hundred years of genetic structure in a sediment revived diatom population. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 4252 (2011). [doi:10.1073/pnas.1013528108](https://doi.org/10.1073/pnas.1013528108) [Medline](#)
30. A. F. Bennett, R. E. Lenski, in *In the Light of Evolution. Volume 1. Adaptation and Complex Design*, J. C. Avise, F. J. Ayala, Eds. (National Academies Press, 2007), pp. 225–238.
31. J. L. Knies, R. Izem, K. L. Supler, J. G. Kingsolver, C. L. Burch, The genetic basis of thermal reaction norm evolution in lab and natural phage populations. *PLoS Biol.* **4**, e201 (2006). [doi:10.1371/journal.pbio.0040201](https://doi.org/10.1371/journal.pbio.0040201) [Medline](#)

32. I. E. Huertas, M. Rouco, V. López-Rodas, E. Costas, Warming will affect phytoplankton differently: evidence through a mechanistic approach. *Proc. Biol. Sci.* **278**, 3534 (2011). [doi:10.1098/rspb.2011.0160](https://doi.org/10.1098/rspb.2011.0160) [Medline](#)
33. S. L. Chown *et al.*, Adapting to climate change: A perspective from evolutionary physiology. *Clim. Res.* **43**, 3 (2010). [doi:10.3354/cr00879](https://doi.org/10.3354/cr00879)
34. M. J. Angilletta Jr., R. S. Wilson, C. A. Navas, R. S. James, Tradeoffs and the evolution of thermal reaction norms. *Trends Ecol. Evol.* **18**, 234 (2003). [doi:10.1016/S0169-5347\(03\)00087-9](https://doi.org/10.1016/S0169-5347(03)00087-9)
35. J. Frantz, P. Novak, g3data (2000) (available at <http://www.frantz.fi/software/g3data.php>).
36. A. Gelman, J. Hill, *Data Analysis Using Regression and Multilevel/Hierarchical Models* (Cambridge University Press, Cambridge, UK, 2007), p. 624.
37. K. P. Burnham, D. R. Anderson, *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach* (Springer-Verlag, New York, ed. 2nd, 2002), p. 488.
38. A. S. Schwaderer *et al.*, Eco-evolutionary differences in light utilization traits and distributions of freshwater phytoplankton. *Limnol. Oceanogr.* **56**, 589 (2011). [doi:10.4319/lo.2011.56.2.0589](https://doi.org/10.4319/lo.2011.56.2.0589)
39. A. J. Kerkhoff, W. F. Fagan, J. J. Elser, B. J. Enquist, Phylogenetic and growth form variation in the scaling of nitrogen and phosphorus in the seed plants. *Am. Nat.* **168**, E103 (2006). [doi:10.1086/507879](https://doi.org/10.1086/507879) [Medline](#)
40. M. D. Guiry, G. M. Guiry, AlgaeBaseWorld-wide electronic publication, National University of Ireland, Galway (2012) (available at <http://www.algaebase.org>).
41. ITIS, Integrated Taxonomic Information System, (2012) (available at <http://www.itis.gov>).
42. J. E. Bissinger, D. J. S. Montagnes, J. Sharples, D. Atkinson, Predicting marine phytoplankton maximum growth rates from temperature: Improving on the Eppley curve using quantile regression. *Limnol. Oceanogr.* **53**, 487 (2008). [doi:10.4319/lo.2008.53.2.0487](https://doi.org/10.4319/lo.2008.53.2.0487)
43. R. Lande, Natural selection and random genetic drift in phenotypic evolution. *Evolution* **30**, 314 (1976). [doi:10.2307/2407703](https://doi.org/10.2307/2407703)
44. C. J. E. Metcalf *et al.*, Evolution of flowering decisions in a stochastic, density-dependent environment. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 10466 (2008). [doi:10.1073/pnas.0800777105](https://doi.org/10.1073/pnas.0800777105) [Medline](#)
45. D. Z. Childs, T. N. Coulson, J. M. Pemberton, T. H. Clutton-Brock, M. Rees, Predicting trait values and measuring selection in complex life histories: reproductive allocation decisions in Soay sheep. *Ecol. Lett.* **14**, 985 (2011). [doi:10.1111/j.1461-0248.2011.01657.x](https://doi.org/10.1111/j.1461-0248.2011.01657.x) [Medline](#)
46. P. A. Abrams, 'Adaptive Dynamics' vs. 'adaptive dynamics'. *J. Evol. Biol.* **18**, 1162 (2005). [doi:10.1111/j.1420-9101.2004.00843.x](https://doi.org/10.1111/j.1420-9101.2004.00843.x) [Medline](#)
47. S. M. Griffies *et al.*, Formulation of an ocean model for global climate simulations. *Ocean Science* **1**, 45 (2005). [doi:10.5194/os-1-45-2005](https://doi.org/10.5194/os-1-45-2005)

48. R Core Team, (2012). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org/>.
49. B. Bolker and R Development Core Team (2012). *bbmle*: Tools for general maximum likelihood estimation. R package version 1.0.5.2. [http://CRAN.R-project.org/package = bbmle](http://CRAN.R-project.org/package=bbmle)
50. H. Wickham, Reshaping data with the reshape package. *J. Stat. Softw.* **21**, 12 (2007).
51. H. Wickham, *ggplot2: elegant graphics for data analysis*. Springer New York (2009).
52. D. Pierce, *ncdf4*: Interface to Unidata netCDF (version 4 or earlier) format data files. R package version 1.0 (2010) (available at [http://cran.r-project.org/package = ncdf4](http://cran.r-project.org/package=ncdf4)).
53. E. J. Pebesma, Multivariable geostatistics in S: the gstat package. *Comput. Geosci.* **30**, 683 (2004). [doi:10.1016/j.cageo.2004.03.012](https://doi.org/10.1016/j.cageo.2004.03.012)
54. D. Bates, M. Maechler, B. Bolker, *lme4*: Linear mixed-effects models using S4 classes. R package version 0.999375-42 (2011). [http://cran.r-project.org/package = lme4](http://cran.r-project.org/package=lme4).
55. V. Vona *et al.*, Temperature responses of growth, photosynthesis, respiration and NADH: Nitrate reductase in cryophilic and mesophilic algae. *New Phytol.* **163**, 325 (2004). [doi:10.1111/j.1469-8137.2004.01098.x](https://doi.org/10.1111/j.1469-8137.2004.01098.x)
56. M.-L. Teoh, W.-L. Chu, H. Marchant, S.-M. Phang, Influence of culture temperature on the growth, biochemical composition and fatty acid profiles of six Antarctic microalgae. *J. Appl. Phycol.* **2**, 421 (2005).
57. X. Wang, K. W. Tang, Y. Wang, W. O. Smith, Jr., Temperature effects on growth, colony development and carbon partitioning in three *Phaeocystis* species. *Aquat. Biol.* **9**, 239 (2010). [doi:10.3354/ab00256](https://doi.org/10.3354/ab00256)
58. M. Fiala, L. Oriol, Light-temperature interactions on the growth of Antarctic diatoms. *Polar Biol.* **10**, 629 (1990). [doi:10.1007/BF00239374](https://doi.org/10.1007/BF00239374)
59. G. Jacques, Some ecophysiological aspects of the Antarctic phytoplankton. *Polar Biol.* **2**, 27 (1983). [doi:10.1007/BF00258282](https://doi.org/10.1007/BF00258282)
60. L. Rhodes, B. Peake, A. L. MacKenzie, S. Marwick, *Coccolithophores Gephyrocapsa oceanica* and *Emiliana huxleyi* (Prymnesiophyceae = Haptophyceae) in New Zealand's coastal waters: Characteristics of blooms and growth in laboratory culture. *N. Z. J. Mar. Freshw. Res.* **29**, 345 (1995). [doi:10.1080/00288330.1995.9516669](https://doi.org/10.1080/00288330.1995.9516669)
61. E. T. Buitenhuis, T. Pangerc, D. J. Franklin, C. Le Quéré, G. Malin, Growth rates of six coccolithophorid strains as a function of temperature. *Limnol. Oceanogr.* **53**, 1181 (2008). [doi:10.4319/lo.2008.53.3.1181](https://doi.org/10.4319/lo.2008.53.3.1181)
62. M. K. de Boer, E. M. Koolmees, E. G. Vrieling, A. M. Breeman, M. van Rijssel, Temperature responses of three *Fibrocapsa japonica* strains (Raphidophyceae) from different climate regions. *J. Plankton Res.* **27**, 47 (2005). [doi:10.1093/plankt/fbh149](https://doi.org/10.1093/plankt/fbh149)
63. M. H. Conte, A. Thompson, D. Lesley, R. P. Harris, Genetic and physiological influences on the alkenone/alkenoate versus growth temperature relationship in *Emiliana huxleyi* and

- Gephyrocapsa oceanica. *Geochim. Cosmochim. Acta* **62**, 51 (1998). [doi:10.1016/S0016-7037\(97\)00327-X](https://doi.org/10.1016/S0016-7037(97)00327-X)
64. J. A. Cannon, in *Toxic Phytoplankton Blooms in the Sea*, T. J. Smayda, Y. Shimizu, Eds. (Elsevier Science, 1993), pp. 741–745.
65. S. M. Renaud, H. C. Zhou, D. L. Parry, L.-V. Thinh, K. C. Woo, Effect of temperature on the growth, total lipid content and fatty acid composition of recently isolated tropical microalgae *Isochrysis* sp., *Nitzschia closterium*, *Nitzschia paleacea*, and commercial species *Isochrysis* sp. (clone T. ISO). *J. Appl. Phycol.* **7**, 595 (1995). [doi:10.1007/BF00003948](https://doi.org/10.1007/BF00003948)
66. S. M. Renaud, L.-V. Thinh, G. Lambrinidis, D. L. Parry, Effect of temperature on growth, chemical composition and fatty acid composition of tropical Australian microalgae grown in batch cultures. *Aquaculture* **211**, 195 (2002). [doi:10.1016/S0044-8486\(01\)00875-4](https://doi.org/10.1016/S0044-8486(01)00875-4)
67. P. D. Chappell, E. A. Webb, A molecular assessment of the iron stress response in the two phylogenetic clades of *Trichodesmium*. *Environ. Microbiol.* **12**, 13 (2010). [doi:10.1111/j.1462-2920.2009.02026.x](https://doi.org/10.1111/j.1462-2920.2009.02026.x) [Medline](#)
68. K. J. S. Meier, C. Höll, H. Willems, Effect of temperature on culture growth and cyst production in the calcareous dinoflagellates *Calciodinellum albatrosianum*, *Leonella granifera* and *Pernambugia tuberosa*. *Micropaleontology* **50**, (Suppl_1), 93 (2004). [doi:10.2113/50.Suppl_1.93](https://doi.org/10.2113/50.Suppl_1.93)
69. E. M. Hulburt, R. R. L. Guillard, The relationship of the distribution of the diatom *Skeletonema tropicum* to temperature. *Ecology* **49**, 337 (1968). [doi:10.2307/1934464](https://doi.org/10.2307/1934464)
70. E. M. Hulburt, The adaptation of marine phytoplankton species to nutrient and temperature. *Ocean Science and Engineering* **7**, 187 (1982).
71. G. Usup, D. M. Kulis, D. M. Anderson, Growth and toxin production of the toxic dinoflagellate *Pyrodinium bahamense* var. *compressum* in laboratory cultures. *Nat. Toxins* **2**, 254 (1994). [doi:10.1002/nt.2620020503](https://doi.org/10.1002/nt.2620020503) [Medline](#)
72. W. H. Thomas, Effects of temperature and illuminance on cell division rates of three species of tropical oceanic phytoplankton. *J. Phycol.* **2**, 17 (1966). [doi:10.1111/j.1529-8817.1966.tb04586.x](https://doi.org/10.1111/j.1529-8817.1966.tb04586.x)
73. L. I. Falcón, S. Pluvinaige, E. J. Carpenter, Growth kinetics of marine unicellular N₂-fixing cyanobacterial isolates in continuous culture in relation to phosphorus and temperature. *Mar. Ecol. Prog. Ser.* **285**, 3 (2005). [doi:10.3354/meps285003](https://doi.org/10.3354/meps285003)
74. H. K. Schöne, The influence of light and temperature on the growth rates of six phytoplankton species from the upwelling area off Northwest Africa. *Rapports et Procès-verbaux des Réunion* **180**, 246 (1982).
75. T. G. Chin, C. F. Chen, S. C. Liu, S. S. Wu, Influence of temperature and salinity on the growth of three species of planktonic diatoms. *Oceanol. Limnol. Sin.* **7**, 373 (1965).
76. J. Bollmann, C. Klaas, L. E. Brand, Morphological and physiological characteristics of *Gephyrocapsa oceanica* var. *typica* Kamptner 1943 in culture experiments: evidence for genotypic variability. *Protist* **161**, 78 (2010). [doi:10.1016/j.protis.2009.08.002](https://doi.org/10.1016/j.protis.2009.08.002) [Medline](#)

77. C. J. Band-Schmidt, Effects of growth medium, temperature, salinity and seawater source on the growth of *Gymnodinium catenatum* (Dinophyceae) from Bahia Concepcion, Gulf of California, Mexico. *J. Plankton Res.* **26**, 1459 (2004). [doi:10.1093/plankt/fbh133](https://doi.org/10.1093/plankt/fbh133)
78. L. McKay, D. Kamykowski, E. Milligan, B. Schaeffer, G. Sinclair, Comparison of swimming speed and photophysiological responses to different external conditions among three *Karenia brevis* strains. *Harmful Algae* **5**, 623 (2006). [doi:10.1016/j.hal.2005.12.001](https://doi.org/10.1016/j.hal.2005.12.001)
79. R. W. Krawiec, Autecology and clonal variability of the marine centric diatom *Thalassiosira rotula* (Bacillariophyceae) in response to light, temperature and salinity. *Mar. Biol.* **69**, 79 (1982). [doi:10.1007/BF00396964](https://doi.org/10.1007/BF00396964)
80. L. R. Moore, R. Goericke, S. W. Chisholm, Comparative physiology of *Synechococcus* and *Prochlorococcus*: Influence of light and temperature on growth, pigments, fluorescence and absorptive properties. *Mar. Ecol. Prog. Ser.* **116**, 259 (1995). [doi:10.3354/meps116259](https://doi.org/10.3354/meps116259)
81. C. M. James, S. Al-Hinty, A. E. Salman, Growth and ω 3 fatty acid and amino acid composition of microalgae under different temperature regimes. *Aquaculture* **77**, 337 (1989). [doi:10.1016/0044-8486\(89\)90218-4](https://doi.org/10.1016/0044-8486(89)90218-4)
82. S. Kahn, O. Arakawa, Y. Onoue, Physiological investigations of a neurotoxin-producing phytoflagellate, *Chattonella marina* (Raphidophyceae). *Aquacult. Res.* **29**, 9 (1998).
83. N. Watabe, K. M. Wilbur, Effects of temperature on growth, calcification, and coccolith form in *Coccolithus huxleyi*. *Limnol. Oceanogr.* **11**, 567 (1966). [doi:10.4319/lo.1966.11.4.0567](https://doi.org/10.4319/lo.1966.11.4.0567)
84. W. S. Maddux, R. F. Jones, Some interactions of temperature, light intensity, and nutrient concentration during the continuous culture of *Nitzschia closterium* and *Tetraselmis* sp. *Limnol. Oceanogr.* **9**, 79 (1964). [doi:10.4319/lo.1964.9.1.0079](https://doi.org/10.4319/lo.1964.9.1.0079)
85. W. H. Thomas, A. N. Dodson, C. A. Linden, Optimum light and temperature requirements for *Gymnodinium splendens*, a larval fish food organism. *Fish Bull.* **71**, 599 (1973).
86. T. Yamatogi, M. Sakaguti, N. Takagi, M. Iwataki, K. Matsuoka, Effects of temperature, salinity and light intensity on the growth of a harmful dinoflagellate *Cochlodinium polykrikoides* Margalef occurring in coast waters of West Kyushu, Japan. *Bull. Plankton Soc. Japan* **52**, 4 (2005).
87. M. Yamaguchi, T. Honjo, Effects of temperature, salinity and irradiance on the growth of the noxious red tide flagellate *Gymnodinium nagasakiense* (Dinophyceae). *Nippon Suisan Gakkai Shi* **55**, 2029 (1989). [doi:10.2331/suisan.55.2029](https://doi.org/10.2331/suisan.55.2029)
88. R. L. Miller, D. L. Kamykowski, Effects of temperature, salinity, irradiance and diurnal periodicity on growth and photosynthesis in the diatom *Nitzschia americana*: Light-saturated growth. *J. Phycol.* **22**, 339 (1986). [doi:10.1111/j.1529-8817.1986.tb00033.x](https://doi.org/10.1111/j.1529-8817.1986.tb00033.x)
89. Y. Nakamura, M. M. Watanabe, Growth characteristics of *Chattonella antiqua* (Raphidophyceae). *J. Oceanogr. Soc. Jpn* **39**, 110 (1983). [doi:10.1007/BF02070796](https://doi.org/10.1007/BF02070796)
90. T. Nishikawa, M. Yamaguchi, Effect of temperature on light-limited growth of the harmful diatom *Eucampia zodiacus* Ehrenberg, a causative organism in the discoloration of *Porphyra thalli*. *Harmful Algae* **5**, 141 (2006). [doi:10.1016/j.hal.2005.06.007](https://doi.org/10.1016/j.hal.2005.06.007)

91. K. Ono, S. Khan, Y. Onoue, Effects of temperature and light intensity on the growth and toxicity of *Heterosigma akashiwo* (Raphidophyceae). *Aquacult. Res.* **31**, 427 (2000). [doi:10.1046/j.1365-2109.2000.00463.x](https://doi.org/10.1046/j.1365-2109.2000.00463.x)
92. E. Breitbarth, A. Oschlies, J. LaRoche, Physiological constraints on the global distribution of *Trichodesmium* – effect of temperature on diazotrophy. *Biogeosciences* **4**, 53 (2007). [doi:10.5194/bg-4-53-2007](https://doi.org/10.5194/bg-4-53-2007)
93. S. H. Baek, S. Shimode, T. Kikuchi, Growth of dinoflagellates, *Ceratium furca* and *Ceratium fusus* in Sagami Bay, Japan: The role of temperature, light intensity and photoperiod. *Harmful Algae* **7**, 163 (2008). [doi:10.1016/j.hal.2007.06.006](https://doi.org/10.1016/j.hal.2007.06.006)
94. Y. Suzuki, M. Takahashi, Growth responses of several diatom species isolated from various environments to temperature. *J. Phycol.* **31**, 880 (1995). [doi:10.1111/j.0022-3646.1995.00880.x](https://doi.org/10.1111/j.0022-3646.1995.00880.x)
95. H. A. Barker, The culture and physiology of the marine dinoflagellates. *Arch. Mikrobiol.* **6**, 157 (1935). [doi:10.1007/BF00407285](https://doi.org/10.1007/BF00407285)
96. R. R. L. Guillard, J. H. Ryther, Studies of marine planktonic diatoms. I. *Cyclotella nana* Hustedt, and *Detonula confervacea* (Cleve) Gran. *Can. J. Microbiol.* **8**, 229 (1962). [doi:10.1139/m62-029](https://doi.org/10.1139/m62-029) [Medline](#)
97. J. H. Ryther, The ecology of phytoplankton blooms in Moriches Bay and Great South Bay, Long Island, New York. *Biol. Bull.* **106**, 198 (1954). [doi:10.2307/1538713](https://doi.org/10.2307/1538713)
98. P. A. Thompson, M.-X. Guo, P. J. Harrison, Effects of variation in temperature. I. On the biochemical composition of eight species of marine phytoplankton. *J. Phycol.* **28**, 481 (1992). [doi:10.1111/j.0022-3646.1992.00481.x](https://doi.org/10.1111/j.0022-3646.1992.00481.x)
99. H. Curl Jr., G. C. McLeod, The physiological ecology of a marine diatom, *Skeletonema costatum* (Grev.) Cleve. *J. Mar. Res.* **19**, 70 (1961).
100. D. Karentz, T. J. Smayda, Temperature and seasonal occurrence patterns of 30 dominant phytoplankton species in Narragansett Bay over a 22-year period (1959-1980). *Mar. Ecol. Prog. Ser.* **18**, 277 (1984). [doi:10.3354/meps018277](https://doi.org/10.3354/meps018277)
101. T. J. Smayda, Experimental observations on the influence of temperature, light, and salinity on cell division of the marine diatom, *Detonula confervacea* (Cleve) Gran. *J. Phycol.* **5**, 150 (1969). [doi:10.1111/j.1529-8817.1969.tb02596.x](https://doi.org/10.1111/j.1529-8817.1969.tb02596.x)
102. C. R. Tomas, *Olisthodiscus luteus* (Chrysophyceae) I. Effects of salinity and temperature on growth, motility and survival. *J. Phycol.* **14**, 309 (1978). [doi:10.1111/j.1529-8817.1978.tb00303.x](https://doi.org/10.1111/j.1529-8817.1978.tb00303.x)
103. L. Ignatiades, T. J. Smayda, Autecological studies on the marine diatom *Rhizosolenia fragilissima* Bergon. I. The influence of light, temperature, and salinity. *J. Phycol.* **6**, 332 (1970).
104. P. G. Falkowski, The adenylate energy charge in marine phytoplankton: The effect of temperature on the physiological state of *Skeletonema costatum* (Grev.) Cleve. *J. Exp. Mar. Biol. Ecol.* **27**, 37 (1977). [doi:10.1016/0022-0981\(77\)90052-1](https://doi.org/10.1016/0022-0981(77)90052-1)

105. E. G. Durbin, Studies on the autecology of the marine diatom *Thalassiosira nordenskioldii* Cleve. I. The influence of daylength, light intensity, and temperature on growth. *J. Phycol.* **10**, 220 (1974).
106. M. W. Fawley, Effects of light intensity and temperature interactions on growth characteristics of *Phaeodactylum tricornutum* (Bacillariophyceae). *J. Phycol.* **20**, 67 (1984). [doi:10.1111/j.0022-3646.1984.00067.x](https://doi.org/10.1111/j.0022-3646.1984.00067.x)
107. C. J. Watras, S. W. Chisholm, D. M. Anderson, Regulation of growth in an estuarine clone of *Gonyaulax tamarensis* Lebour: Salinity-dependent temperature responses. *J. Exp. Mar. Biol. Ecol.* **62**, 25 (1982). [doi:10.1016/0022-0981\(82\)90214-3](https://doi.org/10.1016/0022-0981(82)90214-3)
108. I. Bravo, D. M. Anderson, The effects of temperature, growth medium and darkness on excystment and growth of the toxic dinoflagellate *Gymnodinium catenatum* from northwest Spain. *J. Plankton Res.* **16**, 513 (1994). [doi:10.1093/plankt/16.5.513](https://doi.org/10.1093/plankt/16.5.513)
109. D. Grzebyk, B. Berland, Influences of temperature, salinity and irradiance on growth of *Prorocentrum minimum* (Dinophyceae) from the Mediterranean Sea. *J. Plankton Res.* **18**, 1837 (1996). [doi:10.1093/plankt/18.10.1837m](https://doi.org/10.1093/plankt/18.10.1837m)
110. S. M. Etheridge, C. S. Roesler, Effects of temperature, irradiance, and salinity on photosynthesis, growth rates, total toxicity, and toxin composition for *Alexandrium fundyense* isolates from the Gulf of Maine and Bay of Fundy. *Deep Sea Res. Part II Top. Stud. Oceanogr.* **52**, 2491 (2005). [doi:10.1016/j.dsr2.2005.06.026](https://doi.org/10.1016/j.dsr2.2005.06.026)
111. R. El-Sabaawi, P. J. Harrison, Interactive effects of irradiance and temperature on the photosynthetic physiology of the pennate diatom *Pseudo-nitzschia granii* (Bacillariophyceae) from the northeast subarctic Pacific. *J. Phycol.* **42**, 778 (2006). [doi:10.1111/j.1529-8817.2006.00246.x](https://doi.org/10.1111/j.1529-8817.2006.00246.x)
112. N. I. Lewis, S. S. Bates, J. L. McLachlan, J. C. Smith, in *Toxic Phytoplankton Blooms in the Sea*, T. J. Smayda, Y. Shimizu, Eds. (Elsevier Science, 1993), pp. 601–606.
113. G. Kräbs, C. Büchel, Temperature and salinity tolerances of geographically separated *Phaeodactylum tricornutum* Böhlin strains: maximum quantum yield of primary photochemistry, pigmentation, proline content and growth. *Bot. Mar.* **54**, 231 (2011). [doi:10.1515/bot.2011.037](https://doi.org/10.1515/bot.2011.037)
114. A. M. Johnston, The effect of environmental variables on ¹³C discrimination by two marine phytoplankton. *Mar. Ecol. Prog. Ser.* **132**, 257 (1996). [doi:10.3354/meps132257](https://doi.org/10.3354/meps132257)
115. S. Khan, O. Arakawa, Y. Onoue, Growth characteristics of a neurotoxin-producing chloromonad *Fibrocapsa japonica* (Raphidophyceae). *J. World Aquacult. Soc.* **27**, 247 (1996). [doi:10.1111/j.1749-7345.1996.tb00606.x](https://doi.org/10.1111/j.1749-7345.1996.tb00606.x)
116. L. A. Hobson, Effects of interactions of irradiance, daylength, and temperature on division rates of three species of marine unicellular algae. *J. Fish. Res. Board Can.* **31**, 391 (1974). [doi:10.1139/f74-066](https://doi.org/10.1139/f74-066)
117. N. Grimm, T. Weisse, Die temperaturabhängigkeit des wachstums von *Phaeocystis pouchetii* (Haptophyceae) in batchkulturen. *Helgol. Meeresunters.* **39**, 201 (1985). [doi:10.1007/BF01997450](https://doi.org/10.1007/BF01997450)

118. M. Ø. Jensen, Ø. Moestrup, Autecology of the toxic dinoflagellate *Alexandrium ostenfeldii*: Life history and growth at different temperatures and salinities. *Eur. J. Phycol.* **32**, 9 (1997). [doi:10.1080/09541449710001719325](https://doi.org/10.1080/09541449710001719325)
119. B. Edvardsen, E. Paasche, Two motile stages of *Chrysochromulina polylepis* (Prymnesiophyceae): Morphology, growth and toxicity. *J. Phycol.* **28**, 104 (1992). [doi:10.1111/j.0022-3646.1992.00104.x](https://doi.org/10.1111/j.0022-3646.1992.00104.x)
120. A. M. Sundström *et al.*, *Gymnodinium corollarium* sp. nov. (Dinophyceae) - A new cold-water dinoflagellate responsible for cyst sedimentation events in the Baltic Sea. *J. Phycol.* **45**, 938 (2009). [doi:10.1111/j.1529-8817.2009.00712.x](https://doi.org/10.1111/j.1529-8817.2009.00712.x)
121. J. Thronsen, Occurrence and productivity of small marine flagellates. *Nord. J. Bot.* **23**, 269 (1976).
122. M. V. Nielsen, Growth and chemical composition of the toxic dinoflagellate *Gymnodinium galatheanum* in relation to irradiance, temperature and salinity. *Mar. Ecol. Prog. Ser.* **136**, 205 (1996). [doi:10.3354/meps136205](https://doi.org/10.3354/meps136205)
123. M. V. Nielsen, C. P. Tønseth, Temperature and salinity effect on growth and chemical composition of *Gyrodinium aureolum* Hulbert in culture. *J. Plankton Res.* **13**, 389 (1991). [doi:10.1093/plankt/13.2.389](https://doi.org/10.1093/plankt/13.2.389)
124. A. Larsen, S. Bryant, Growth rate and toxicity of *Prymnesium parvum* and *Prymnesium patelliferum* (Haptophyta) in response to changes in salinity, light and temperature. *Sarsia* **83**, 409 (1998).
125. D. S. Reay, D. B. Nedwell, J. Priddle, J. C. Ellis-Evans, Temperature dependence of inorganic nitrogen uptake: reduced affinity for nitrate at suboptimal temperatures in both algae and bacteria. *Appl. Environ. Microbiol.* **65**, 2577 (1999). [Medline](https://pubmed.ncbi.nlm.nih.gov/10501100/)
126. E. Nordli, Experimental studies on the ecology of *Ceratia*. *Oikos* **8**, 200 (1957). [doi:10.2307/3564999](https://doi.org/10.2307/3564999)
127. M. Montesor, C. R. Tomas, Growth and probably gamete formation in the marine dinoflagellate *Ceratium schrankii*. *J. Phycol.* **24**, 495 (1988).
128. G. Mjaaland, Some laboratory experiments on the coccolithophorid *Coccolithus huxleyi*. *Oikos* **7**, 251 (1956). [doi:10.2307/3564925](https://doi.org/10.2307/3564925)
129. C. Lovejoy *et al.*, Distribution, phylogeny, and growth of cold-adapted picoprasinophytes in Arctic seas. *J. Phycol.* **43**, 78 (2007). [doi:10.1111/j.1529-8817.2006.00310.x](https://doi.org/10.1111/j.1529-8817.2006.00310.x)
130. Z. I. Johnson *et al.*, Niche partitioning among *Prochlorococcus* ecotypes along ocean-scale environmental gradients. *Science* **311**, 1737 (2006). [doi:10.1126/science.1118052](https://doi.org/10.1126/science.1118052)
[Medline](https://pubmed.ncbi.nlm.nih.gov/16411000/)
131. A. R. Juhl, Growth rates and elemental composition of *Alexandrium monilatum*, a red-tide dinoflagellate. *Harmful Algae* **4**, 287 (2005). [doi:10.1016/j.hal.2004.05.003](https://doi.org/10.1016/j.hal.2004.05.003)
132. E. R. Zinser *et al.*, Influence of light and temperature on *Prochlorococcus* ecotype distributions in the Atlantic Ocean. *Limnol. Oceanogr.* **52**, 2205 (2007). [doi:10.4319/lo.2007.52.5.2205](https://doi.org/10.4319/lo.2007.52.5.2205)

133. B. J. Binder, D. M. Anderson, Physiological and environmental control of germination in *Scripsiella trochoidea* (Dinophyceae) resting cysts. *J. Phycol.* **23**, 99 (1987).
134. H. Kaeriyama *et al.*, Effects of temperature and irradiance on growth of strains belonging to seven *Skeletonema* species isolated from Dokai Bay, southern Japan. *Eur. J. Phycol.* **46**, 113 (2011). [doi:10.1080/09670262.2011.565128](https://doi.org/10.1080/09670262.2011.565128)
135. J. Navarro, M. Munoz, A. Contreras, Temperature as a factor regulating growth and toxin content in the dinoflagellate *Alexandrium catenella*. *Harmful Algae* **5**, 762 (2006). [doi:10.1016/j.hal.2006.04.001](https://doi.org/10.1016/j.hal.2006.04.001)